

Review Article

Targeting the Ras signaling pathway: a rational, mechanism-based treatment for hematologic malignancies?

Christoph W. M. Reuter, Michael A. Morgan, and Lothar Bergmann

A series of alterations in the cellular genome affecting the expression or function of genes controlling cell growth and differentiation is considered to be the main cause of cancer. These mutational events include activation of oncogenes and inactivation of tumor suppressor genes. The elucidation of human cancer at the molecular level allows the design of rational, mechanism-based therapeutic agents that antagonize the specific activity of biochemical processes that are essential to the malignant phenotype of cancer cells. Because the frequency of *RAS* mutations is among the highest for any gene in human cancers, development of inhibitors of the Ras-mitogen-activated pro-

tein kinase pathway as potential anticancer agents is a very promising pharmacologic strategy. Inhibitors of Ras signaling have been shown to revert Ras-dependent transformation and cause regression of Ras-dependent tumors in animal models. The most promising new class of these potential cancer therapeutics are the farnesyltransferase inhibitors. The development of these compounds has been driven by the observation that oncogenic Ras function is dependent upon posttranslational modification, which enables membrane binding. In contrast to many conventional chemotherapeutics, farnesyltransferase inhibitors are remarkably specific and have been demon-

strated to cause no gross systemic toxicity in animals. Some orally bioavailable inhibitors are presently being evaluated in phase II clinical trials. This review presents an overview on some inhibitors of the Ras signaling pathway, including their specificity and effectiveness in vivo. Because Ras signaling plays a crucial role in the pathogenesis of some hematologic malignancies, the potential therapeutic usefulness of these inhibitors is discussed. (Blood. 2000;96:1655-1669)

© 2000 by The American Society of Hematology

The *RAS* gene family

At the cellular surface, many different receptors are expressed that allow cellular response to extracellular signals provided by the environment. After ligand binding, receptor activation leads to a large variety of biochemical events in which small guanosine triphosphate hydrolases (GTPases; eg, Ras) are crucial. Ras proteins are prototypical G-proteins that have been shown to play a key role in signal transduction, proliferation, and malignant transformation. G-proteins are a superfamily of regulatory GTP hydrolases that cycle between 2 conformations induced by the binding of either guanosine diphosphate (GDP) or GTP¹⁻³ (Figure 1). The Ras-like small GTPases are a superfamily of proteins that include Ras, Rap1, Rap2, R-Ras, TC21, Ral, Rheb, and M-Ras. The *RAS* gene family consists of 3 functional genes, *H-RAS*, *N-RAS*, and *K-RAS*. The *RAS* genes encode 21-kd proteins, which are associated with the inner leaflet of the plasma membrane (H-Ras, N-Ras, and the alternatively spliced K-RasA and K-RasB). Whereas H-Ras, N-Ras, and K-RasB are ubiquitously expressed, K-RasA is induced during differentiation of pluripotent embryonal stem cells in vitro.⁴

Regulatory proteins that control the GTP/GDP cycling rate of Ras include GTPase-activating proteins (GAPs), which accelerate the rate of GTP hydrolysis to GDP, and guanine nucleotide exchange factors (GEFs; eg, SOS and CDC25), which induce the dissociation of GDP to allow association of GTP.³ In the GTP-bound state, Ras couples the signals of activated growth factor

receptors to downstream mitogenic effectors. By definition, proteins that interact with the active GTP-bound form of Ras (and thus become GTP-dependently activated) to transmit signals are called Ras effectors.⁵⁻⁸ Mechanisms by which GTP-Ras influences the activity of its effectors include direct activation (eg, B-Raf, PI-3 kinase), recruitment to the plasma membrane (eg, c-Raf-1), and association with substrates (eg, Ral-GDS). Other candidates for Ras effectors include protein kinases, lipid kinases, and GEFs.^{3,5-8}

Posttranslational modification of Ras

Ras proteins are produced as cytoplasmatic precursor proteins and require several posttranslational modifications to acquire full biologic activity. These modifications include prenylation, proteolysis, carboxymethylation, and palmitoylation⁹⁻¹³ (Figure 2).

Prenylation of proteins by intermediates of the isoprenoid biosynthetic pathway represents a newly discovered form of posttranslational modification and is catalyzed by 3 different enzymes: protein farnesyltransferase (F'ase), protein geranylgeranyltransferase type I (GGTase I), and geranylgeranyltransferase type II (GGTase II).⁹⁻¹³ Prenylated proteins share characteristic carboxy-terminal consensus sequences and can be separated into the proteins with a CAAX (C, cysteine; A, aliphatic amino acid; X, any amino acid) motif and proteins containing a CC or CXC

From the Department of Internal Medicine III (Hematology and Oncology), University of Ulm, Ulm, Germany.

Submitted August 24, 1999; accepted March 30, 2000.

Supported by a grant to C.W.M.R. from the German Research Council (Deutsche Forschungsgemeinschaft Ro 664/I-1) and a grant from the University of Ulm (P.541).

Reprints: C. Reuter, Dept of Internal Medicine III, University of Ulm, Robert-Koch-Str 8, D-89081 Ulm, Germany; e-mail: christoph.reuter@med3.hi.uni-ulm.de.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 U.S.C. section 1734.

© 2000 by The American Society of Hematology

1656 REUTER et al

BLOOD, 1 SEPTEMBER 2000 • VOLUME 96, NUMBER 5




Figure 1. The switch function of Ras. Ras cycles between the active GTP-bound and the inactive GDP-bound state. Mitogenic signals activate guanine GEFs such as SOS and CDC25. GEFs increase the rate of dissociation of GDP and stabilize the nucleoside-free form of Ras, leading to binding of GTP to Ras proteins. Ras can also be activated by the inhibition of the GAPs.

sequence.¹⁴⁻¹⁷ FTase I transfers a farnesyl group from farnesyl-diphosphate (FPP), and GGase I transfers a geranylgeranyl group from geranylgeranyldiphosphate (GGPP) to the cysteine residue of the CAAX motif.¹⁸ GGase II transfers the geranylgeranyl groups from GGPPs to both cysteine residues of CC or CXC motifs.

Farnesylation is the first step in the posttranslational modification of Ras. This modification occurs by covalent attachment of a 15-carbon farnesyl moiety in a thioether linkage to the carboxy-terminal cysteine of proteins that contain the CAAX motif. The reaction is catalyzed by FTase, a heterodimer consisting of a 48-kd and a 45-kd subunit (α_{FTase} and β_{FTase}). Binding sites for the substrates, FPP and the CAAX motif, are located on the α_{FTase} and β_{FTase} subunits.¹⁹⁻²¹ Substrates for FTase include all known Ras proteins, nuclear lamins A and B, the γ -subunit of the retinal trimeric G-protein transducin, rhodopsin kinase, and a peroxisomal protein termed Pxl.⁹⁻¹³

Farnesylation of Ras proteins is followed by endoproteolytic removal of the 3 carboxy-terminal amino acids (AAX) by a cellular thiol-dependent zinc metalloprotease.²² This endoproteolytic activity (RACE, or Ras and a-factor converting enzyme) is a composite of 2 different CAAX proteases: a zinc-dependent activity encoded




Figure 2. Overview of the posttranslational modifications of Ras proteins in cells. FTase or GGase I transfers a farnesyl (F) group or a geranylgeranyl group from FPP or GGPP to the thiol group of the cysteine residue in the CAAX motif. The C-terminal tripeptide is removed by a CAAX-specific endoprotease in the endoplasmic reticulum. A PPMase attaches the methyl group from S-adenosylmethionine (SAM) to the C-terminal cysteine. Finally, a prenyl protein-specific palmitoyltransferase (PPMTase) attaches palmitoyl groups (P) to cysteines near the farnesylated C-terminus. PPM1I indicates prenyl protein-specific methyltransferase inhibitor.

by AFC1 and the type IIb signal peptidase-like RCE1 (Ras converting enzyme 1).²³ The final step in the carboxy-terminal modification of proteins with a CAAX motif (eg, Ras) is the methylation of the carboxyl group of the prenylated cysteine residue by an as yet uncharacterized methyltransferase.

Some Ras proteins (H-Ras, N-Ras, Ras2) are further lipidated by palmitoylation at 1 or 2 cysteines near the farnesylated carboxy-terminus.^{9-13,24-27} Like farnesylation, H-Ras palmitoylation plays an important role for signaling functions in vivo.²⁷ Microinjection experiments in *Xenopus* oocytes revealed that palmitoylation of H-Ras dramatically enhances its affinity for membranes as well as its ability to activate mitogen-activated protein kinase (MAPK) and initiate meiotic maturation.^{11,27} Both a Ras-specific protein (palmitoyltransferase) and a palmitoyl-protein (thioesterase) have been characterized.^{28,29} In contrast to farnesylation and proteolysis, palmitoylation and methylation of Ras are thought to be reversible and may have a regulatory role.^{11,12}

The Ras-to-MAPK signal transduction pathway

The MAPK signaling cascades

MAPK pathways are well-conserved major signaling systems involved in the transduction of extracellular signals into cellular responses in a variety of organisms, including mammals.³⁰⁻³⁵ The core components of the MAPK signaling cascades are 3 sequential kinases, including MAP kinase (MAPK, or extracellular signal-regulated kinase, ERK), MAPK kinase (MAPKK, or MAPK/ERK kinase, MEK), and MAPKK kinase (MAPKKK, or MEK kinase, MEKK) (Figure 3). The MAPKs are activated by dual phosphorylation on tyrosine and threonine residues by upstream dual-specificity MAPKKs. MAPKKs are also phosphorylated and activated by serine- and threonine-specific MAPKKKs. At least 6 MAPK cascades have been clearly identified in mammalian




Figure 3. The best-characterized MAPK modules are the ERK pathway, the SAPK/JNK pathway, and the p38 pathway. The classical Ras-to-MAPK cascade is shown in bold. The MAPK cascades consist of a MAPKKK, a MAPKK, and a MAPK. MAPKKKs are activated through a large variety of extracellular signals such as growth factors, differentiation factors, and stress. The activated MAPKKK can phosphorylate and activate 1 or several MAPKKs, which, in turn, phosphorylate and activate a specific MAPK. Activated MAPK phosphorylates and activates various substrates in the cytoplasm and the nucleus of the cell, including transcription factors. These downstream targets control cellular responses (eg, apoptosis, proliferation, and differentiation). Thick arrows connect the signaling proteins with their preferred substrates (effectors). Note the complexity and the potential for crosstalk between the pathways.

cells.³⁰⁻³⁵ The best characterized MAPK signaling pathways are (1) the Ras-to-MAPK signal transduction pathway (or ERK pathway), which is responsive to signals from receptor tyrosine kinase, hematopoietic growth factor receptors, and some heterotrimeric G-protein-coupled receptors, which promote cell proliferation or differentiation; (2) the stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) pathway, which is activated in response to stresses such as heat, high osmolarity, UV irradiation, and proinflammatory cytokines such as tumor necrosis factor- α and interleukin-1 (IL-1); and (3) the p38 pathway, which is responsive to osmotic stress, heat shock, lipopolysaccharide, tumor necrosis factor- α , and IL-1 (Figure 3).³⁰⁻³⁵ Scaffold/adaptor proteins such as MP-1, JSAP-1, and JIP-1 route MAPK modules in mammals by binding ERK-1 and MEK-1, JNK-3 and SEK-1 and MEKK-1, or JNK and MKK-7 and MEKs, respectively.^{34,35}

Ras-to-MAPK signaling via receptor tyrosine kinases and cytokine receptors

The Ras-to-MAPK pathway appears to be an essential shared element of mitogenic signaling. The MAPKs ERK-1 and ERK-2 are activated by various mitogens in all cells. Ras functions as a membrane-associated biologic switch that relays signals from ligand-stimulated receptors to cytoplasmic MAPK cascades. These receptors include G-protein-coupled serpentine receptors, tyrosine kinase receptors (eg, platelet-derived growth factor receptor [PDGFR], epidermal growth factor [EGF] receptor) and cytokine receptors that cause stimulation of associated nonreceptor tyrosine kinases (NRTKs; eg, Src, Lyn, Fes). Ligand binding to the extracellular domain of receptor tyrosine kinases (RTKs) causes receptor dimerization, stimulation of protein tyrosine kinase activity, and autophosphorylation.³⁶⁻⁴⁰ Tyrosine autophosphorylation sites in growth factor receptors (eg, EGF receptor) function as high-affinity binding sites for SH-2 (src homology) domains of signaling molecules such as PI-3 kinase (PI-3K), phospholipase C (PLC)- γ , p120-GAP, Shc, and SHP-2 tyrosine phosphatase.³⁹

In contrast to receptor tyrosine kinases, cytokine receptors (such as the prototypical IL-3, IL-5, GM-CSF receptors) do not contain a kinase domain. These receptors are heterodimers of a ligand-specific α -subunit and a β -subunit that is common to IL-3, IL-5, and granulocyte-macrophage colony-stimulating factor (GM-CSF) receptors.^{41,42} The NRTKs Lyn and Fes and the Janus kinase JAK2 are physically associated with the β -subunit. The conserved proline-rich motifs in the α - and β -subunits (eg, IL-3, IL-5, GM-CSF-R, IL-2-R, G-CSF-R, and erythropoietin-R) are critical for JAK2 binding and activation. (Figure 4). After ligand binding and receptor dimerization, the receptor-bound tyrosine kinases become activated and cause a cascade of tyrosine phosphorylations. As in the receptor tyrosine kinases, these phosphotyrosines represent docking sites for many signaling molecules, including adaptor proteins (eg, PI-3K, Shc, SHP-2, Grb-2).^{41,42}

The SH3 domain of Grb-2 binds to SOS, which is a GEF for Ras and facilitates the replacement of GDP with GTP.^{3-8,36-40} When Ras becomes GTP-loaded, Ras effectors (such as Rafs, MEKK, PI-3K, and Ral) bind to Ras and become activated. The Raf kinases (A-Raf, B-Raf, c-Raf-1) are important Ras effectors and have been demonstrated to act as MAPKKs/MEKKs in the Ras-to-MAPK (or ERK) pathway.^{30-40,43-45} Raf kinases have been shown to selectively phosphorylate and activate MAPKKs MEK-1 and MEK-2.^{36-40,43-45} Other MEK-1/MEK-2 activators include TPL-2, MEKK-1, and c-Mos.⁴⁶⁻⁴⁸ MEK-1 and MEK-2 are dual-specificity kinases that activate the MAPKs of the ERK subgroup (ERK-1 and ERK-2).^{30-35,49-52} ERK-1 and ERK-2 are proline-directed protein




Figure 4. The classical Ras-to-MAPK cascade. (A) Signaling by cytokine receptors. The IL-3, IL-6, and GM-CSF receptors consist of a ligand-specific α -subunit and a common β -subunit. The β -subunit binds the NRTKs Lyn, Fes, and JAK2. After ligand binding, the α - and β -subunits are thought to dimerize, thus activating the receptor-bound NRTKs and subsequently causing a cascade of tyrosine phosphorylations. The phosphotyrosine residues represent docking sites for various signaling molecules (eg, Shc, SHP-2). ERKs are activated via the classical Ras-to-MAPK pathway. In addition, the MAPKs p38 and JNK become activated. The activation pathway is not completely understood, but some lines of evidence support the involvement of Ras or HPK-1 (hematopoietic progenitor kinase, a mammalian Ste20-related protein). Activated JAK2 phosphorylates the STAT (signal transducers and activators of transcription) family of nuclear factors which form heterodimers and homodimers, thus causing their translocation to the nucleus and subsequent binding to γ -activating sequences of the promoter region of various genes.^{41,42} (B) Signaling by receptor tyrosine kinases. Extracellular stimuli such as mitogens or stress result in the intracellular activation of different MAPK cascades. The ERK-1/2 pathway is activated by mitogens in all cells and is an essential part of mitogenic signaling. Translocation of a fraction of activated ERKs to the nucleus subsequently leads to activation of transcription factors such as Elk-1, CREB, SRF, and Fos.⁴⁰ The Raf kinases connect upstream tyrosine kinases and Ras with downstream serine/threonine kinases. When Ras becomes GTP-loaded, Rafs bind to Ras. It is unclear if Ras-Raf binding is itself always sufficient to activate the Raf kinases, which subsequently phosphorylate and activate the downstream MEKs. GTP-Ras also binds and activates PI-3K and Ral-GEF. PI-3K produces lipid second messengers, which activate Akt (Akt kinase) and nPKC. Ral-GEF activates Ral-GTPases by promoting the GTP-bound state of Ral. Ral-GTP binds to Ral-BP1 (a GAP for CDC42 and Rac), phospholipase D (PLD1), and Ca^{2+} calmodulin (CaM). (I) Inhibitors of Ras membrane association (eg, FTI, GGT1, PPM1, and REPI); (II) sulindac; (III) MEK inhibitors (eg, PD098059, U0126, and Ro 09-2110). The thick, black arrows show the classical Ras-to-MAPK cascade. The thick, open arrows represent the Ras-to-Ral and the Ras-to-PI-3K signaling pathways. The STAT pathway is shown on the left.

kinases that phosphorylate Ser/Thr-Pro motifs in the consensus sequence Pro-Xaa-Ser/Thr-Pro, where Xaa is any amino acid and $n = 1$ or 2. Several cytoplasmic and nuclear substrates of the ERKs have been identified. The best-characterized ERK substrates are cytoplasmic phospholipase A₂ (cPLA₂), the ribosomal protein S6 kinases (RSKs), and the transcription factor Elk-1.^{30,32,53,54}

The Ras-to-Ral and the Ras-to-PI-3K signaling pathways

Since the discovery of Raf as a direct Ras effector, numerous other putative Ras effectors have been identified.³⁻⁸ Among these, evidence to date best supports "effector" roles for the Ral-GEFs (Ral-GDS, RGL, and RGF) and the p110 subunit of PI-3K.^{3-8,55,56} (Figure 4).

Ral-GEFs are activated via binding to GTP-Ras. Ral-GEFs in turn activate Ral-GTPases by promoting the GTP-bound state of Ral. As members of the Ras subfamily of Ras-related GTPases, Ral proteins (RalA and RalB) also cycle between the active GTP-bound states and inactive GDP-bound states. Ral-GTP binds Ral-BP1 (Ral-binding protein-1 or Rlipl = Rlipl [Ral-interacting protein-1]), which is a GAP for CDC42 and Rac. These 2 GTPases are

Table 1. Activation of Ras in hematologic malignancies

Malignancy	Type of activation	Frequency (%)	Reference nos.
	Point mutations of RAS		
Multiple myeloma	N-, K-RAS	30-40	74, 76-78
Plasma cell leukemia	N-, K-RAS	60-70	74, 76-78
Acute myeloid leukemia (AML)	N-, K-RAS	20-30	60-71
Childhood AML	N-, K-RAS	20-40	64, 72
Acute lymphoblastic leukemia	N-, K-RAS	20	68, 69, 75
Chronic myelomonocytic leukemia (CMML)	N-, K-RAS	50-70	85, 73, 80
Juvenile myelomonocytic myeloid leukemia (JMML)	N-RAS	30	107
	Mutation of c-Kit/c-FMS receptor family		
Myeloproliferative disorder, mastocytosis	c-Kit	10	83-85
AML	FLT-3	20	87
	CSF-1(c-FMS)	10-20	81, 82
	Fusion tyrosine kinases		
Chronic myeloid leukemia	Bcr-Abl, t(9;22), Ras-GTP	95	95-97
CMML	Tet-PDGFR β , t(5;12)		91, 92
AML	Tet-Abl, t(12;9)		93, 94
Anaplastic large cell lymphoma	Npm-AIk, t(2;5)	30-40	89, 90
	Inactivation of tumor suppressors		
JMML	Inactivation of NF-1, (Ras-GAP)		90-108

Ras proteins are small GTPases that cycle between 2 conformations induced by the binding of GDP or GTP. In the active, GTP-bound conformation, Ras binds to and activates effector proteins such as Raf kinases, PI-3K, and Ral-GDS. Mutations in codons 12, 13, or 61 of the RAS genes lead to activated Ras proteins that have lost the ability to become inactivated and thus stimulate growth autonomously. Activated tyrosine receptor kinases, which are upstream of Ras (eg, mutated c-Kit, c-FMS, FLT-3, or activated fusion tyrosine kinases such as BCR-Abl, Tet-Abl, Npm-AIk, and Tet-PDGFR β), may also cause elevated levels of active, GTP-bound Ras and thus stimulate cell proliferation. The loss of the tumor suppressor NF-1, a Ras-GTPase activating protein (Ras-GAP), also causes Ras activation.

involved in the regulation of the actin cytoskeleton, the SAPK/JNK pathway, and the p38 pathway (Figure 3).

Ras-GTP also binds to and activates the catalytic domain of PI-3K. The lipid second-messenger molecules produced (eg, phosphatidylinositol phosphates PtdIns 3,4-P₂ and PtdIns 3,4,5-P₃) activate the phosphoinositide-dependent kinases PDK-1 and PDK-2, which then activate Akt kinase and nonconventional isoforms of protein kinase C (ncPKC). PI-3K has been implicated in 4 apparently distinct cellular functions, including mitogenic signaling (DNA synthesis), inhibition of apoptosis, intracellular vesicle trafficking and secretion, and regulation of actin and integrin functions. These functions are most likely mediated by distinct phosphoinositide products of PI-3K⁵⁶ (Figure 4).

Role of Ras activation in hematologic malignancies

The constitutive activation of Ras appears to be an important factor for the malignant growth of human cancer cells. Recently, the Ras-related proteins R-Ras, M-Ras, and TC21 have also been shown to possess transforming activities similar to those of Ras.⁵⁷⁻⁵⁹ However, their role in human malignancies is unclear. Mutations of the RAS proto-oncogenes (H-RAS, N-RAS, K-RAS) are frequent genetic aberrations found in 20% to 30% of all human tumors, although the incidences in tumor type vary greatly.^{60,61} The highest rate of RAS mutations was detected in adenocarcinomas of the pancreas (90%), the colon (50%), and the lung (30%). In follicular and undifferentiated carcinomas of the thyroid, the incidence of RAS mutations is also considerable (50%). The most commonly observed RAS mutations arise at sites critical for Ras regulation—namely, codons 12, 13, and 61. Each of these mutations results in the abrogation of the normal GTPase activity of Ras. While all the Ras mutants still form complexes with GAP, the GTPase reaction of Ras cannot be stimulated by GAP, thus causing an increase in the half-lives of Ras-GTP mutants.^{1,5} Transformation

results, at least in part, from unregulated stimulation of the mitogenic signal transduction pathway.^{60,61}

Ras activation is frequently observed in hematologic malignancies such as myeloid leukemias and multiple myelomas. In about one-third of the myelodysplastic syndromes (MDS) and acute myeloid leukemias (AML), RAS genes are mutationally activated⁶²⁻⁷³ (Table 1). N-RAS is mutated and activated in most of the cases, and the presence of the mutation is not associated with any particular FAB type, cytogenetic abnormality, or clinical feature, including prognosis.⁷⁴ RAS mutations occur in about 40% of newly diagnosed multiple myeloma patients, and the frequency increases with disease progression.⁷⁴ Mutations in N-RAS—especially codon 61 mutations—are more frequent than K-RAS mutations.⁷⁴⁻⁷⁸

In addition to activation by mutation, Ras is thought to be deregulated by constitutive activation of proto-oncogenes and inactivation of tumor suppressor genes.^{79,80} Several types of human cancers show oncogenic activation of RTKs or NRTKs. Constitutively activated versions of normal receptor tyrosine kinases contain single point mutations (eg, CSF-1 receptor, the Neu/erbB-2 receptor, and the c-Kit receptor), duplications of juxtamembrane domain-coding sequences (eg, FLT3 receptor), or deletions of the negative regulatory regions in the ligand binding or the transmembrane domains (eg, ErbB-3 receptor). Point mutations of the CSF-1 receptor (c-FMS) at codons 301 and 969 were found in 10% to 20% of AML or MDS.^{81,82} Point mutations in the catalytic domain of the c-Kit receptor are found in some cases of myeloproliferative disorders and in 10% of the patients with mastocytosis.⁸³⁻⁸⁵ Furthermore, activating tandem internal duplication of the FLT3 receptor has been reported in 20% of AML.⁸⁶ The members of the c-Kit/c-FMS receptor kinase family (eg, c-Kit, c-FMS, FLT3) are linked with components of the Ras-to-MAPK signaling pathway (eg, Grb-2 and Shc), suggesting that activating mutations of c-FMS and FLT3 may induce activation of Ras.^{87,88}

In addition, translocations involving receptor tyrosine kinases produce chimeric proteins in which varying N-terminal portions of either the ligand-binding or the transmembrane domain are replaced

with novel protein sequences.^{79,80} Several of these chimeric proteins have been found in human hematologic malignancies. The Npm-Alk fusion protein, a fusion of the N-terminal portion of Npm with the entire cytoplasmic domain of the receptor tyrosine kinase Alk, is generated by the t(2;5) chromosomal translocation in anaplastic large cell lymphoma.^{80,81} Tel-PDGFR β is a fusion protein consisting of the transcription factor Tel (translocation, Ets, leukemia) and PDGFR β , a well-known receptor tyrosine kinase.^{91,92} It is generated by the t(5;12) translocation in a subset of chronic myelomonocytic leukemias that results in receptor dimerization and activation and thus leads to the constitutive activation of the Ras-MAPK pathway.³ Another Tel fusion protein, Tel-Abl, is generated by the t(12;9) translocation in AML.^{93,94} Abl is an NRTK that is also mutated and activated in chronic myelogenous leukemia.⁹⁵⁻⁹⁷ In Bcr-Abl, the product of the t(9;22) translocation, the N-terminal Bcr portion serves as an oligomerization domain. Bcr-Abl is a constitutively activated cytosolic tyrosine kinase that causes abrogation of growth factor dependence, blockade of differentiation, and direct inhibition of apoptosis. Although Ras mutations are extremely rare in chronic myelogenous leukemia, the involvement of Ras has been demonstrated in Bcr-Abl⁺ cells by the presence of increased levels of GTP-Ras, which leads to the activation of the Raf kinases and other Ras effectors.⁹⁵⁻⁹⁷ Thus, the deregulation of Ras function appears to be a common theme in the transformation by activated receptor and NRTKs. Ras activation may cause elevated cell cycle progression and inhibition of apoptosis.^{71,79,80,95-97}

In addition to oncogenes, tumor suppressor genes have also been found to be involved in the deregulation of Ras. Neurofibromin, the product of the NF1 gene, encodes a Ras-GAP and is mutated in the autosomal dominant type 1 neurofibromatosis.⁹⁸ Interestingly, neurofibromatosis type 1 is associated with an increased tendency to develop myeloid leukemias, especially juvenile myelomonocytic myeloid leukemia (JMML).⁹⁹⁻¹⁰⁷ About 15% of children with JMML cases have clinical neurofibromatosis.⁹⁹ Additionally, inactivating mutations of the NF1 gene have been found in 15% of JMML without clinical diagnosis of neurofibromatosis, suggesting the existence of NF1 mutations in approximately 30% of all JMML cases.^{100,102} The involvement of Ras is demonstrated by the finding that leukemic cells from children with neurofibromatosis type 1 show a moderate elevation in the percentage of GTP-Ras.¹⁰⁰⁻¹⁰⁶ Furthermore, 15% to 30% of JMML cases lacking the NF1 mutation have activating RAS mutations.¹⁰⁷ The observation that human JMML cells exhibit hypersensitivity to GM-CSF suggests a common pathophysiologic mechanism involving downstream Ras signaling.¹⁰⁶⁻¹⁰⁸

The pathophysiologic importance of the Ras-MAPK signaling pathway is underscored by the positioning of several oncogene and tumor suppressor gene products on this pathway (Figure 4). Furthermore, it has recently been demonstrated that mutant N-RAS induces myeloproliferative disorders resembling human chronic myelogenous leukemia, AML, and apoptotic syndromes similar to human MDS in bone marrow-repopulated mice.¹⁰⁹ These observations make Ras and the Ras-MAPK pathway an attractive target for the development of new anticancer agents.

Inhibitors of the Ras-MAPK pathway

Inhibitors of Ras farnesyltransferase

Elimination of Ras function by homologous gene recombination or antisense RNA has demonstrated that expression of activated Ras is necessary for maintaining the transformed phenotype of tumor cells.¹¹⁰⁻¹¹³ Inhibitors of oncogenic Ras activity may therefore

prove useful as anticancer agents against Ras-induced tumors. One strategy to impede oncogenic Ras function *in vivo* is the inhibition of Ras posttranslational modification. It has been demonstrated that mutation of the evolutionarily conserved CAAX motif in Ras abolishes plasma membrane binding as well as transforming activity.¹¹⁴⁻¹²¹ Although Ras undergoes several steps of posttranslational modification, only farnesylation is necessary for its membrane localization and cell-transforming activity.¹²¹ Therefore, it has been proposed that the activity of oncogenic Ras could be blocked by inhibiting the FTase responsible for this modification. However, many CAAX-containing proteins need additional palmitoylation for stable membrane association.

FTase has become a very attractive target for the development of anticancer agents because control of Ras farnesylation can control the function of oncogenic Ras.¹¹⁴⁻¹²¹ Numerous inhibitors of Ras FTase have been synthesized or identified. These Ras FTase inhibitors can be grouped into 3 classes: (1) FPP analogues such as (α -hydroxyfarnesyl) phosphonic acid, β -ketophosphonic and β -hydroxyphosphonic acid derivatives, and J-104871¹²²⁻¹²⁴ (Figure 5A); (2) CAAX peptide analogues such as BZA-5B, BZA-2B,¹²⁵⁻¹²⁷ L-731,734, L-731,735, L-739,749,¹²⁸⁻¹³² L-739,787,¹³³ L-739,750, L-744,832,^{129,134-137} B581,¹³⁸ Cys-4-ABA-Met and Cys-AMBA-Met,¹³⁹ FTI-276, FTI-277,¹⁴⁰⁻¹⁴³ B956, and its methyl ester B1096¹⁴⁴ (Figure 5B); in addition, nonpeptidic, tricyclic FTase inhibitors have been developed such as SCH144342, SCH54429, SCH159228, and SCH66336¹⁴⁵⁻¹⁴⁹ (Figure 6); and (3) bisubstrate inhibitors such as phosphonic acid analogues, the phosphinate inhibitors BMS-185878 and BMS-186511, the phosphonate inhibitor BMS-184467, phosphinyl acid-based derivatives, and the hydroxamine acid analogues¹⁵⁰⁻¹⁵² (Figure 5C).

In addition to chemically synthesized compounds, several natural products have been identified as FTase inhibitors. These include limonene,¹⁵³ manumycin (UCF1-C) and its related compounds UCF1-A and UCF1-B,¹⁵⁴⁻¹⁵⁶ chaetomelic acid A and B, zaragozic acids, peptidocinnamins, gliotoxin,¹¹⁵ barceloneic acid A,¹⁵⁷ RPR113228,¹⁵⁸ actinoplanic acids A and B,¹⁵⁹ oreganic acid,¹⁶⁰ lupane derivatives,¹⁶¹ saquayamycins,¹⁶² valinotcin A and its analogues,¹⁶³ and ganoderic acid A and C.¹⁶⁴

Effects of FTase inhibitors in intact tumor cells. Several FTase inhibitors were demonstrated to be active in intact cells (Table 2). These compounds have been shown to modulate several critical aspects of Ras transformation in whole cells, including selective inhibition of anchorage-independent growth of Ras-transformed fibroblasts in soft agar, morphologic reversion of the Ras-induced phenotype, and inhibition of oocyte maturation induced by oncogenic Ras without gross cytotoxic effects on normal cells. One of the first FTase inhibitors found to be active in intact tumor cells, the FPP analogue (α -hydroxyfarnesyl) phosphonic acid, only partially inhibited the farnesylation of Ras in H-Ras-transformed NIH3T3 fibroblasts.¹⁶⁵ Subsequently, more potent FTase inhibitors have been developed. L-739,749 inhibited growth of Ras-transformed rat fibroblasts and caused rapid morphologic reversion of the transformed phenotype.¹³⁰ The compound B581 inhibited colony formation of Ras-transformed cells in soft agar.¹³⁸ BZA-5B and BZA-2B, both benzodiazepine peptidomimetic FTase inhibitors, have been shown to slow the growth of H-Ras-transformed cells at concentrations that do not affect the growth of nontransformed cells.^{125-127,166,167} The peptidomimetic FTase inhibitor B956 and its methyl ester B1086 inhibited the formation of soft agar colonies of 14 human tumor cell lines expressing oncogenic forms of H-Ras, N-Ras, and K-Ras.¹⁴⁴ Five human tumor cell lines expressing




Figure 5. Structures of FPP-, CAAX-based, and bisubstrate inhibitors of Ftsase. (A) Chemical structures of FPP and FPP-based inhibitors of Ftsase and PPMtsase. FPP is composed of a hydrophobic farnesyl group and a highly charged pyrophosphate moiety. The basic structural element in the Ftsase inhibitors is a farnesyl group, a pyrophosphate isostere, and a linker. (B) CAAX-based Ftsase inhibitors. Structural comparison between CAAX-based Ftsase inhibitors of the pseudopeptide class and the CAAX tetrapeptides ClfM and CvfM. The potent, nonsubstrate Ftsase inhibitors ClfM and CvfM were identified by systematic amino acid replacements within the CAAX sequence. In F11-276 and F11-277, the AA residues of the CAAX motif have been replaced by a hydrophobic linker. (C) In bisubstrate Ftsase inhibitors, the farnesyl group of FPP and the tripeptide group of the CAAX motif are connected via a linker.

wild-type Ras required higher concentrations of the drug to inhibit colony formation. About 50% of K-Ras-transformed cell lines were observed to be as resistant as non-Ras-transformed cell lines. It has been suggested that nontransformed cells may produce a form of Ras that is isoprenylated even in the presence of FTase inhibitors.¹⁴⁴ Additionally, this phenomenon may be due to functional redundancy within the *RAS* family. The tricyclic inhibitor SCH44342 specifically blocks morphologic transformation in-




Figure 6. Nonpeptidic, tricyclic FTase inhibitors. FTase inhibitor SCH44342 had no *in vivo* efficacy. Further substitutions led to SCH66336, a highly potent FTase inhibitor, which was found to have therapeutically useful serum levels and half-lives when given orally to rodents and primates. SCH66336 is being tested in human clinical phase II trials.

duced by Val12-Ha-Ras-CVLS but not Val12-Ha-Ras-CVLI, a form of Ras engineered to bind to GGTase I, indicating that the compound is a specific inhibitor of H-Ras modification by Fase rather than K-Ras modification by GGTase.¹⁴⁵

Similarly, several bisubstrate inhibitors of FTase were shown to inhibit oncogenic Ras-induced growth in vivo. The phosphinyl acid-based bisubstrate analogic FTase inhibitors Nos. 17 to 19 were found to inhibit the anchorage-independent colony growth of Ha-RAS-transformed NIH3T3 cells.¹⁵⁰⁻¹⁵² The bisubstrate FTase inhibitor BMS-186511 inhibited FTase activity in whole cells and produced strong inhibition of Ras-transformed growth. Although both H-Ras- and K-Ras-transformed cells were affected by BMS-186511, K-Ras cells appeared to be less sensitive.¹⁵¹ BMS-186511 did not produce any signs of gross, unspecific cytotoxicity in untransformed normal cells.

The Ffase inhibitor L-744,832 blocked the anchorage-dependent and -independent growth of 31 of 42 human tumor cell lines.¹³⁶ The origin of the tumor cell and the presence or absence of mutationally activated Ras did not correlate with the response to the Ffase inhibitor. Interestingly, cell lines with wild-type Ras and activated receptor tyrosine kinases were also sensitive to L-744,832. In contrast, nontransformed epithelial cell lines were far less sensitive. Recently, L-739,749 and L-744,832 have also been reported to inhibit the colony growth of juvenile myelomonocytic leukemia cells, which are known to exhibit deregulated cytokine signal transduction involving the Ras pathway.¹³⁷

Biologic mechanisms of FfTase inhibitors in intact cells. Recent investigations into the biologic mechanism of the growth inhibition of Ras-transformed cells have shown that farnesylation of K-Ras and N-Ras is more resistant to FfTase inhibitors than farnesylation of H-Ras.^{126,144,167,168} In part, this phenomenon is a result of a 10- to 50-fold higher affinity of FfTase for K-Ras4B than for other Ras isoforms.^{126,170} In the absence of FfTase inhibitors, all Ras proteins are present only in the farnesylated form. However, K-Ras and N-Ras (but not H-Ras) become geranylgeranylated by GGTase I in vivo in a dose-dependent manner when intracellular farnesylation is inhibited by an FfTase inhibitor.^{126,169-171} Subsequently, both FfTase and GGTase I inhibitors are required for inhibition of K-Ras processing.^{168,172} The lack of growth inhibition and gross cytotoxic effects of FfTase inhibitors on normal cells is thought to be a result of the resistance of K-Ras processing to FfTase inhibitors.¹⁶⁷

Treatment of Ras-transformed cells with FTase inhibitors results in selective suppression of Ras-dependent oncogenic signaling. This includes the inhibition of Ras processing, which results in a decrease in the relative amount of fully processed Ras; the progressive, dose-dependent cytoplasmic accumulation of unprocessed Ras and inactive Ras-Raf complexes; inhibition of the Ras-induced constitutive activation of MAPK^{138,140,141,146,173}; and decreased transcriptional activity of both c-Jun and Elk-1.¹³⁸ Transformation by mutationally activated Raf, MEK, Mos, or Fos (all of which are downstream effectors of Ras) is not blocked by FTase inhibitors.^{129,136}

Although FTase inhibitors block Ras farnesylation and the Ras-induced transformed phenotype, proteins other than Ras may be targets of these compounds.^{174,175} Ffase inhibitors block anchorage-independent growth of many human tumor cell lines in soft agar culture, but there is no correlation between biologic susceptibility and the presence of Ras mutations.^{136,144} In addition, anchorage-independent growth of K-Ras-transformed cells is abrogated

Table 2. Effect of FTase inhibitors on intact cells

Compound	Cell type	RAS mutation	Ras activation	IC ₅₀ μ mol/L	Reference nos.
L-731,734	Fibroblasts	v-RAS	+	1	128
L-730,749	Rat-1 fibroblasts		+	2.5	130
	JMML cells		+	1-10	167
B581	NIH3T3 cells	t-RAS	+	—	132
BZA-5B	Rat-1 cells	H-RAS	+	—	138
BZA-2B					125, 126
B956, B1086	Human tumor cell lines	H-RAS	+	0.2-0.7	166, 167
		N-RAS	+	3-7	144
		K-RAS	+	1.7-50	
SCH44342	Cos cells	H-RAS	+	—	145
	Various human tumor cell lines	H-, N-, K-RAS	+/-	1	138
L-744,832	JMML cells		+	1-10	132
Compound no. 46	NIH3T3	H-RAS	+	0.19	187
FTI-277	Glioma cell lines			2.5-15.5	142
Compound no. 10	Rat-1 cells	v-RAS	+	2.5-5	185
Compound 5m	NIH3T3 cells	H-RAS-F	+	0.18	186
FTS*	Rat-1 cells	N-RAS	+	30-40*	108
	Human melanoma cell lines	N-RAS	+/-	100*	

Several FTase inhibitors have been demonstrated to revert specifically the Ras-transformed phenotype and anchorage-independent growth in fibroblasts and human tumor cell lines. Cell growth inhibition may be a result of induction of apoptosis or arrest in the G1 phase of the cell cycle.

† Indicates positive for activated Ras; —, negative for activated Ras.

*FTS, S-farnesylthiosalicylic acid, is an inhibitor of PPMTase.

by FTase inhibitors even though K-Ras processing is not affected.¹⁷² Although it is unclear whether soluble species of oncogenic Ras exert any biologically significant effect in drug-treated cells, it has recently been shown that nonfarnesylated H-Ras proteins can be palmitoylated and thus are biologically active. These proteins bound modestly to the plasma membranes (40%) but were still able to trigger exaggerated differentiation of PC12 cells and potent transformation of NIH3T3 fibroblasts.¹⁷⁶

Recently, it has been suggested that the antitransforming effects of FTase inhibitors are mediated at least in part by alteration of farnesylated Rho proteins, including RhoB.^{174,175,177,178} In contrast to Ras proteins, RhoB exists normally *in vivo* in a farnesylated (RhoB-FF) and a geranylgeranylated version (RhoB-GG).¹⁷⁹ RhoB-GG is essential for the degradation of p27KIP1 and facilitates the progression of cells from G1 to S phase. Treatment with FTase inhibitors results in a loss of RhoB-FF and a gain of RhoB-GG.¹⁷⁸ Expression of a mutant RhoB-GG protein induces phenotypic reversion, cell growth inhibition, and activation of the cell cycle kinase inhibitor p21WAF1 in cells sensitive to FTase inhibitors, including Ras-transformed cells.^{178,180} p21WAF1 mediates the inhibition of cyclinE-associated protein kinase activity, pRB hypophosphorylation, and inhibition of DNA replication, which results in G1 arrest.¹⁸⁰ In addition to the induction of the G1 block, treatment of tumor cells with FTase inhibitors induces apoptosis by upregulating Bax and Bcl-x_s expression and by activating caspases.^{131,181-183}

Synergy of FTase inhibitors with established anticancer treatments such as radiation and chemotherapeutic treatment was recently reported. Agents that prevent microtubule depolymerization, such as taxol and epothilones, act synergistically with FTase inhibitors to block cell growth. FTase inhibitors cause increased sensitivity to induction of the metaphase block by taxol and epothilones.¹⁸⁴ In addition, FTase inhibitors have been shown to increase the radiosensitivity of human tumor cells with activating mutations of RAS oncogenes.¹⁴³

Effects of FTase inhibitors in animal models. FTase inhibitors have also been shown to inhibit the growth of Ras-induced tumors in mouse xenograft models and, more dramatically, in transgenic mouse models (Table 3). Manumycin was reported to inhibit the growth of K-Ras-transformed fibrosarcoma transplanted into nude mice by approximately 70% compared with untreated controls.¹⁵⁴ The CAAX peptide analogue L-739,749 specifically suppressed the tumor growth of H-Ras-, N-Ras-, and K-Ras-induced Rat-1 cell tumors in nude mice by 51% to 66%.¹²⁹ Interestingly, L-739,749 exhibited no evidence of systemic toxicity. The peptidomimetic FTase inhibitors B956 and B1086 were shown to inhibit tumor growth of EJ-1 human bladder carcinoma, HT 1080 human fibrosarcoma and, to a lesser extent, HCT116 human colon carcinoma xenografts in nude mice. Inhibition of Ras processing correlated with the inhibition of the tumor growth by B956.¹⁴⁴ Analogues of the tetrapeptide CVFM,¹⁸⁹ the compound Nos. 46 and 51, showed inhibition of anchorage-independent growth of stably H-Ras-transformed NIH3T3 fibroblasts as well as antitumor activity in an athymic mouse model implanted with H-Ras-transformed Rat-1 cells.¹⁸⁷ J-104871, an FPP-competitive FTase inhibitor, suppressed tumor growth in nude mice transplanted with activated H-RAS-transformed NIH3T3 cells.¹²⁴ In contrast to these results, however, treatment of irradiated mice engrafted with NF-1 deficient hematopoietic cells (-/-) with the FTase inhibitor (FTI) L-744,832 failed to revert a myeloproliferative disorder similar to JMML.¹⁹⁰ Although L-744,832 abrogated the GM-CSF-induced growth, H-Ras processing and MAPK activation of NF-1 (-/-) hematopoietic cells, this FTI did not reduce the constitutively elevated MAPK activity levels in these cells. This may be due to the resistance of N-Ras and K-Ras processing to inhibition by the FTI.¹⁹⁰

In addition to the mouse xenograft models, FTase inhibitors have been tested in transgenic mouse models. The CAAX-based FTase inhibitor L-744,832 induced regression of mammary and salivary carcinomas in MMTV-v-Ha-RAS mice. These mice harbor the viral Ha-RAS oncogene under the control of the mouse

1662 REUTER et al

BLOOD, 1 SEPTEMBER 2000 • VOLUME 96, NUMBER 5

Table 3. Effect of FTase inhibitors in animal models

Compound	Model/tumor	RAS mutation/activation	Dose (mg/kg/d)	Growth inhibition	Reference nos.
Manumycin	Nude mice/K-Ras fibrosarcoma	K-RAS/+		70%	154
	Balb/c nude mice/pancreas (MIA PaCa-2) carcinoma	K-RAS/+	1, 2, 5		191
L-739,749	Nude Harlan mice/Rat-1 cell tumors	H-RAS/+	20	51%-66%	129
		N-RAS/+			
		K-RAS/+			
B956, B1088	Nude mice/bladder (EJ-1) fibrosarcoma (HT1080) colon (HCT 116)	H-RAS/+	100		144
		N-RAS/+			
		K-RAS/+			
L-744,832	MMTV-v-H-RAS transgenic mice/mammary, salivary carcinoma	v-H-RAS/+	10-40	-5.4 vs 16.7	135
			40	-7.7 vs 11.8	134
				-9.9 vs 33.3	
				-12.3 vs 26.3	
FTI-276	Nude Harlan Sprague-Dawley mice/lung (A-549, Calu-1) tumors	K-RAS/+	10, 50, 100	75%	140
FTI-277	NIH3T3 cells	H-RAS-F/+	50	80%	
FTI-276	A/J mice lung adenomas		50	58%	200
FTI-276	Nude Harlan Sprague-Dawley mice/lung (A-549, Calu-1) tumors	K-RAS/+	70	70%-94%	168
GGTI-297*				56%-70%*	
Compound no. 48	Allymic Balb/c	H-RAS/+	45	T/C† = 154%	187
Compound no. 51	Rat-1 cells		45	T/C† = 142%	
Compound 83b, 85b	Nude mice/colon (DLD-1, SW-260) H-Ras-CVLS fibroblasts H-Ras-CVLL fibroblasts	K-RAS/+	10, 50	72%	148
				95%	
				50%	
				100%	
SCH66336	Nude mice/colon (DLD-1, HCT 116) pancreas (MIA PaCa-2) NIH3T3	K-RAS/+	2.5, 10, 40	76%	147
				75%	
				100%	
				100%	
Compound no. 4	Nude mice/colon (DLD-1)	K-RAS/+	10, 50	70%	148
SCH66336	Nude mice/lung (A549, HTB177) pancreas (AsPC-1, HPAF-II) Hs 700 T, MIA PaCa-2) colon (HCT116, DLD-1) prostate (DU-145) urinary bladder (EJ) wap-H-RAS transgenic mice/mammary, salivary tumors	K-RAS/+	2.5, 10, 40	70%, 83%	192
		K-RAS/+		72%, 67%	
		K-RAS/+		67%, 78%	
		K-RAS/+		84%, 76%	
		no mutation		86%	
		H-RAS/+		100%	
		H-RAS/+		67%-86%	
		H-RAS/+			
SCH59228	Allymic mice/colon (DLD-1) H-Ras and K-Ras fibroblast tumors	K-RAS/+	10, 50	>90%	149
L-744,832	MMTV-N-RAS TM transgenic mice/lymphoid and mammary tumors	H-RAS/+	40	-0.7 vs 28.3	137
		N-RAS/+			
L-744,832	MMTV-TGF α /neu transgenic mice/mammary tumors	-/+	40	-7.4 vs 19	183
Compound 5m	Nude mice/NIH3T3	H-RAS-F	150	88	186
FTS*	SCID mice/melanoma (518A2, 607B)	N-RAS	5*	82-90	188

Several FTase inhibitors have shown in vivo antitumor activity in mice. These inhibitors have been demonstrated to cause regression of tumors that depend on activated Ras in mouse xenograft and transgenic mouse models. The growth inhibition is given in percent of controls or as the comparison of the tumor mean growth rate (in mm³/day) in the presence or absence of the FTase inhibitor. Cell growth inhibition may be a result of induction of apoptosis or arrest in the G1 phase of the cell cycle.

*FTS, 8-fomesylthiosalicylic acid, is an inhibitor of PPMTase.

†T/C indicates relative median survival time of treated (T) versus control (C) groups (% T/C values). The activity criterion for increased lifespan was a T/C of $\geq 125\%$.

mammary tumor virus (MMTV) long terminal repeat and develop spontaneous mammary and salivary carcinomas.¹³⁵ In agreement with earlier observations, no systemic toxicity was observed in these mice. Furthermore, L-744,832 was also effective in mammary and lymphoid tumors overexpressing N-RAS in MMTV transgenic mice.¹³⁷ In contrast to H-Ras, N-Ras remained mostly processed. Consistent with these findings, the antineoplastic effect was less intense in the N-RAS model than the H-RAS model.^{135,137} This observation suggests that proteins in addition to Ras may be

targets of the compound. More recently, L-744,832 was shown to induce regression of mammary tumors in MMTV-TGF α and MMTV-TGF α /neu transgenic mice.¹⁸³ Because the mammary tumor cells harbor an activated receptor tyrosine kinase but wild-type Ras, a feature common in breast cancer, these mice provide a useful model system for breast cancer research. Tumor regression by L-744,832 was demonstrated biochemically by inhibition of MAPK activity and biologically by an increase in G1-phase, a decrease in S-phase fractions, and induction of apoptosis.¹⁸³

In both cell culture and mouse models, there is essentially no cytotoxicity or apparent systemic toxicity at doses capable of reverting Ras-induced transformation or of causing tumor regression. FTase inhibitors seem to selectively target a unique aspect of the transformed cell physiology.

Mechanisms of resistance to FTase inhibitors. As with any drug, the development of tumor resistance to FTase inhibitors is an important issue. To date, the relative frequency, the mechanisms, and the development of tumor resistance to FTI are unclear. *K-RAS*-transformed cell lines have been shown to be more resistant to FTase inhibitors than *H-RAS*- or *N-RAS*-transformed cells.^{126,144,167} This phenomenon is thought to be a result of a higher affinity of FTase for K-Ras than for other Ras isoforms.^{126,167} In addition, K-Ras and N-Ras become geranylgeranylated in the presence of FTI.^{169,171} Subsequently, both FTI and geranylgeranyltransferase inhibitor (GGTI) are required for inhibition of K-Ras processing.^{168,172} Recently, a variant *RAS*-transformed cell line was identified that was resistant to phenotypic reversion by FTI.¹⁹³ This phenomenon was not due to mutation of the FTase subunits, changes in intracellular drug accumulation, or amplification of the multiple-drug resistance gene. The precise mechanism of resistance in these cells remained unclear. However, mutational alteration of FTase might also lead to resistance toward FTI. The Y361L mutant of FTase has been shown to exhibit increased resistance to FTI while maintaining FTase activity toward substrates possessing C18 carboxy-termini.¹⁹⁴ Withdrawal of FTI from successfully treated tumor-bearing mice led to subsequent tumor growth in the absence of the drug. A second FTI treatment resulted in a second response in some mice, but some tumors were found to become resistant to FTI.¹³⁵ Therefore, chronic, uninterrupted treatment with FTI might be required.

Inhibitors of geranylgeranyl transferase I

Until recently, the emphasis has been on designing specific FTase inhibitors to block Ras processing. This strategy was employed to avoid possible toxic effects originating from inhibition of GGase I. Because *K-RAS* mutations are most common in human cancers,^{60,61} a critical goal is the development of inhibitors that block the growth of human tumors that harbor K-Ras. The resistance of K-Ras to FTase inhibitors,¹⁶⁷ the lack of potency of FTase inhibitors against K-Ras-transformed cells,¹⁴⁴ and the observation that K-Ras becomes geranylgeranylated in the presence of FTase inhibitors^{126,169-172} led to the development of GGase I inhibitors (Figure 7). GGTI-279, GGTI-287, GGTI-297, and GGTI-298 are CAAL-based peptidomimetics that are selective for GGase I over FTase.^{173,195-199} In contrast, FTI-276 and FTI-277 are CAAM-based peptidomimetics that are potent and selective inhibitors of FTase over GGase I.¹⁷³ H-Ras processing in human tumor cell lines was highly sensitive to FTI-277 and resistant to GGTI-286, whereas K-Ras4B processing was more sensitive to GGTI-286 than FTI-277.¹⁷³ Processing of H-Ras and N-Ras was inhibited by FTI-277, but inhibition of K-Ras processing required both FTase and GGase I inhibitors. Whereas FTI-277 preferentially blocks activation of MAPK by oncogenic H-Ras, GGase inhibitors selectively inhibit the activation of MAPK by oncogenic K-Ras4B.¹⁷³ Although GGTI-298 had very little effect on soft agar growth of several human tumor cell lines harboring H-RAS, N-RAS, or K-RAS mutations, the combination of FTI-277 and GGTI-298 resulted in significant soft agar growth inhibition.¹⁷² Both FTase inhibitors and GGase inhibitors have been reported to arrest Ras-transformed

cells in G0/G1 phase of the cell cycle and to induce apoptosis.^{142,180,196,198,199} In nude mouse xenografts, the GGase inhibitor GGTI-297 suppressed human lung A-549 and Calu-1 carcinoma tumor growth by 60%. However, both FTase and GGase inhibitors were required to inhibit K-Ras processing.¹⁶⁸ Treatment of cells with GGTI-298 blocks PDGF- and EGF-dependent tyrosine phosphorylation of their respective receptors and induces G0/G1-phase arrest and apoptosis.¹⁹⁶⁻¹⁹⁸ GGTI-298 has also been shown to induce the cyclin-dependent kinase inhibitor p21WAF but not p27KIP.¹⁹⁹

Inhibitors of the prenylated protein methyltransferase

The C-terminal prenylated protein methyltransferase (PPMTase) is another potential therapeutically relevant target in the development of inhibitors against the posttranslational processing of Ras. N-acetyl-*trans*, *trans*-farnesyl-L-cysteine (AFC) is a substrate for PPMase and acts as a competitive inhibitor.²⁰¹ Although AFC has been shown to inhibit Ras methylation in Ras-transformed NIH3T3 fibroblasts, it does not inhibit the growth of these cells.²⁰¹ New farnesyl derivatives of rigid carboxylic acid, eg, *S-trans*, *trans*-farnesylthiosalicylic acid (FTS), were found to inhibit the growth of H-Ras-transformed cells and to reverse their transformed morphology by a mechanism unrelated to the inhibition of Ras methylation by PPMase.^{202,203} (Figure 5). It is thought that FTS specifically interacts with Ras farnesylcysteine binding domains and affects membrane anchorage of Ras.^{202,203} In addition, it has been reported that FTS dislodges Ras from H-Ras-transformed cell membranes and renders the Ras protein susceptible to proteolytic degradation.¹⁸⁸ At the same concentration, growth and morphology of non-Ras-transformed or nontransformed cells were not affected by FTS.²⁰³ Despite the lack of FTS-induced




Figure 7. CAAL-based inhibitors of GGase I. GGase I catalyzes the geranylgeranylation of proteins terminating with CAAX sequences where X is restricted to leucine, isoleucine or, to a lesser extent, phenylalanine. In cells, geranylgeranylation of proteins is far more common than farnesylation. Proteins modified by GGase I include Rap1A, Rap1B, Rac1, Rac2, G25K, and RhoA.

cytotoxicity in nontransformed cells, FTS reduced Ras levels in cell membranes and inhibited Ras-dependent cell growth.²⁰³ In contrast to FTase inhibitors (eg, BZA-5B), FTS also inhibited the growth signaling of receptor tyrosine kinases.²⁰³ FTS was shown to decrease total cellular Ras levels, MAPK activity, Raf-1 activity, and DNA synthesis in Ras-transformed EJ-1 cells. This inhibition was also demonstrated in serum-, EGF-, and thrombin-stimulated, untransformed Rat-1 cells.^{204,205} S-farnesyl-thioacetate (FTA), another competitive inhibitor of PPMase, has been shown to suppress growth and induce apoptosis in HL-60 cells.²⁰⁶ Five-chloro- and 4- or 5-fluoro-derivatives of FTS and a C20 S-geranylgeranyl derivative of thiosalicylic acid also cause inhibition of Ras-dependent MAPK activity, DNA synthesis, and EJ-1 cell growth. However, several other derivatives were inactive, suggesting stringent structural requirements for the anti-Ras activity of S-prenyl analogues.²⁰⁷ Recently, FTS was shown (1) to reduce the amount of activated N-Ras and wild-type Ras isoforms in human melanoma cells and Rat-1 fibroblasts, (2) to disrupt ERK signaling, (3) to revert their transformed phenotype, and (4) to cause a significant reduction in the growth of human melanoma in SCID mice.^{188,205}

The dorrigoins are novel antifungal antibiotics that were found to reverse the morphology of Ras-transformed NIH3T3 fibroblasts. Dorrigoins A did not inhibit protein prenylation or protein synthesis but was instead found to inhibit the C-terminal methylation in K-Ras-transformed cells.²⁰⁸

Selective Inhibitors of Ras C-terminal sequence-specific endoprotease

UM96001, TPCK, and BFCCMK are Ras C-terminal sequence-specific endoprotease inhibitors (REPI) and potently inhibit Ras-transformed rat kidney cell growth as well as growth of human cancer cells.²⁰⁹ These compounds have been reported to almost completely block the anchorage-independent clonogenic growth of these cancer cells. REPIs may selectively induce apoptosis in these cells.²⁰⁹

Selective Inhibitors of MAPKKs, or MEK

PD098059 is a synthetic inhibitor of the Ras-MAPK pathway that selectively blocks the activation of MEK-1 and, to a lesser extent, the activation of MEK-2.^{210,211} The inhibition of MEK-1 activation was demonstrated to prevent activation of MAPKs ERK-1/2 and subsequent phosphorylation of MAPK substrates both in vitro and in intact cells. In contrast to FTase inhibitors, PD098059 inhibited stimulation of cell growth by several growth factors.^{210,211} Furthermore, PD098059 reversed the transformed phenotype of Ras-transformed BALB3T3 mouse fibroblasts and rat kidney cells.²¹¹ PD098059 failed to inhibit the stress, and IL-1 stimulated JNK/SAPK and the p38 pathways,²¹⁰ demonstrating its specificity for the ERK pathway. PD098059 has subsequently been used as a tool to study MAPK signaling in various cell types and in carcinogenesis.

Recently, 2 novel inhibitors of MEK-1 and MEK-2 have been identified: U0126^{212,213} and Ro 09-2210.²¹⁴ Ro 09-2210, which was identified by screening microbial broths, exhibits potent antiproliferative effects on activated T cells.²¹⁴ Similarly, U0126 was found to inhibit T-cell proliferation in response to both antigenic stimulation and cross-linked anti-CD3 plus anti-CD28 antibodies.²¹² U0126 and PD098059 are noncompetitive inhibitors with respect to both MEK substrates (ATP and ERK) and bind to free MEK as well as MEK*ERK and MEK*ATP complexes. U0126 displays

significantly higher affinity for all forms of MEK (44- to 357-fold) than does PD098059. U0126 and Ro 09-2210 have an inhibitory concentration of 50% (IC₅₀) of 50 to 70 nmol/L, whereas PD098059 has an IC₅₀ of 5 μ mol/L.²¹²⁻²¹⁴ PD098059 and U0126 impede the growth of Ras-transformed cells in soft agar but show minimal effects on cell growth under normal culture conditions.^{210,213} In contrast to U0126 and PD098059, Ro 09-2210 is also able to inhibit other dual-specificity kinases such as MKK-4, MKK-6, and MKK-7, albeit at 4- to 10-fold higher IC₅₀ concentrations compared with its effect on MEK-1.²¹⁴

Inhibitors of Ras transformation with unknown mechanisms of action

Screening tests for drugs that revert RAS-transformed cells to a normal phenotype led to the identification of a number of compounds, such as azatyrosine, oxanosine, and antipain.²¹⁵⁻²¹⁷ The mechanism by which these compounds revert the RAS-induced phenotype is not understood. The pyrazolo-quinoline compound SCH51344 was identified based on its ability to depress human smooth muscle α -actin promoter activity in RAS-transformed cells. Treatment of *v-abl*-, *v-mos*-, *v-ras*-, *RAS*-, and mutant active MEK-transformed NIH3T3 cells resulted in growth inhibition of these cells in soft agar.²¹⁸ SCH51344 had very little effect on the activities of proteins in the ERK pathway. The ability of SCH51344 to inhibit the anchorage-independent growth of RAC-V12-transformed Rat-1 cells suggests that the point of inhibition is downstream from RAC.²¹⁹

The nonsteroidal, anti-inflammatory drug sulindac has been demonstrated to attenuate the growth and progression of colonic neoplasms in animal models and in patients with familial adenomatous polyposis.^{220,221} Recently, it has been shown that sulindac sulfide (the active metabolite of sulindac) inhibits Ras signaling and transformation by noncovalent binding to the Ras protein. Furthermore, it has been demonstrated that sulindac sulfide impairs Ras-Raf binding, Raf activation, and nucleotide exchange on Ras and that it accelerates the Ras-GTPase reaction.²²² Sulindac is being investigated in a randomized study for the prevention of colon cancer (protocol RUH-SSH-190-0698, NCI-V98-1425).

Disruption of the Ras-to-MAPK signaling pathway has also been shown for the benzoquinone ansamycin geldanamycin. Geldanamycin binds to HSP90 and disrupts the HSP90-Raf-1 multimolecular complex, which causes destabilization of Raf-1 through enhanced degradation of Raf-1.²²³ However, the geldanamycin-HSP90 complex also causes depletion of other HSP90 substrates such as protein kinases and nuclear hormone receptors (including mutant p53 and ErbB2).²²⁴ Several National Cancer Institute-sponsored clinical phase I trials are currently studying the effects of geldanamycin analogues in patients with advanced malignancies.

Conclusions and future directions

FTase and GGTase inhibitors have strong growth inhibitory and antitumor activity in cell culture and animal tumor models without showing nonspecific gross toxicity in animals. The specificity and the lack of nonspecific toxicity contrasts dramatically with the nonspecificity and high toxicity of currently available chemotherapeutic drugs. The recent development of orally bioavailable FTase inhibitors with potent and selective in vivo antitumor activity underscores their potential usefulness in the future treatment of human malignancies. The observation that FTase and GGTase

Table 4. FTase inhibitors in clinical trials

Compound	Phase	Malignancy	Status	Protocol ID
R115777	I	Solid advanced tumors	Completed	NCI-97-C-0086B*
R115777	I	Refractory solid tumors (children)	Active	NCI-98-C-0141*
R115777	I	Refractory or recurrent acute leukemia or accelerated or blastic phase chronic myeloid leukemia	Active	MSGCC-9802 NCI-T99-0030* MSGCC-0398115 URCC-980300
R115777	I	Advanced cancer	Active	UTHSC-9785011335 NCI-V98-1501 JRF-R115777 SACI-IDO-98-03
L-778,123	I	Refractory or recurrent solid tumors or lymphomas	Active	MSKCC-98116 NCI-G99-1572* MERCK-003-04
SCH66336	I	Advanced cancer	Active	MSKCC-99020 NCI-G99-1540* SPRI-C98-564-01
SCH66336	II	Metastatic adenocarcinoma of the pancreas	Active	MSKCC-98115 NCI-G99-1571* SPRI-C98-545-12
SCH66336	II	Metastatic adenocarcinoma of the pancreas	Active	CWRU-SCH-1298 NCI-G99-1534* SPRI-C98-545-18
SCH66336	II	Metastatic adenocarcinoma of the pancreas refractory to gemcitabine	Not yet active	UCLA-9906030 NCI-G99-1610* SPRI-P00346
SCH66336	II	Previously treated, inoperable, locally advanced, or metastatic transitional cell carcinoma of the bladder	Not yet active	CAN-NCIC-IND128 SPRI-CAN-NCIC-IND128

Three orally bioavailable FTase inhibitors have entered several phase I/II clinical trials. Most of these trials are National Cancer Institute-sponsored (*). The malignancies comprise a wide variety of human solid tumors (eg, primary brain tumors such as gliomas, neurofibromas, neurofibrosarcomas and malignant schwannomas, neuroblastomas, Wilms tumors, germ cell tumors, adenocarcinomas of the pancreas) and hematopoietic diseases such as acute leukemias and chronic myeloid leukemia in acceleration and blastic phase.

inhibitors induce apoptosis in treated tumor cells as well as a G0-G1 arrest suggests that they are not merely cytostatic but cytotoxic for tumor cells. However, the absence of toxicity due to FTase inhibitors in normal cells and tissues in mice at doses that inhibit tumor growth is poorly understood. Ras knockout experiments have demonstrated that H-RAS- and N-RAS-deficient mice are born and grow normally, whereas K-RAS-deficient embryos die between embryonic day 12.5 and term. This finding suggests a partial functional overlap within the RAS gene family.²²⁵⁻²²⁸ However, H-RAS and N-RAS cannot compensate for the loss of K-RAS function in K-RAS deficient mice. Functionally redundant pathways might allow normal cells to tolerate treatment with FTase inhibitors.

Because mutated RAS genes have a high prevalence in human cancers (eg, pancreatic, lung, and colon cancers), inhibitors specific for FTase, GGTase, and MEK were initially designed to block the Ras-to-MAPK signaling in solid tumor cells. More than 90% of RAS mutations found in human tumors occur in N-RAS or K-RAS. Whereas the reversion of the H-RAS-induced transformation by FTase inhibitors correlates well with the intracellular inhibition of H-Ras processing, N-Ras and K-Ras are cross-prenylated by GGTase I in cells treated with FTase inhibitors. However, many of these N-RAS- or K-RAS-transformed cell lines (and even tumor cell lines that do not harbor RAS mutations) are sensitive to FTase inhibitors. Cell biology studies suggest that FTase and GGTase inhibitors may act at additional levels beyond the inhibition of Ras processing. The exact mechanism of action has emerged as a question of major interest, especially because transformed tumor cells respond to treatment with these inhibitors while normal cells

remain largely unaffected. Non-Ras targets of FTase and GGTase inhibitors may include other cellular proteins (eg, Rho) that are farnesylated or geranylgeranylated.^{174,175,178,229-231}

FTase inhibitors (eg, R115777, L-778,123, and SCH66336) have entered several phase I/II clinical trials (Table 4). These trials are still ongoing, and preliminary results have not been published. Because favorable synergistic effects have been described for combinations of FTase inhibitors with traditional anticancer treatments such as radiation and chemotherapy,^{143,184} it will be interesting to see if these results translate into improved patient outcome in clinical trials. The high prevalence of mutationally activated Ras in solid tumors has been the driving force of Ras inhibitor research. However, recent studies in cell culture and animal models suggest that transformed cells with an activated Ras pathway (eg, via mutations upstream of Ras) are also highly sensitive for FTase inhibitors. The involvement of N-RAS in the molecular pathophysiology of myeloid leukemias and multiple myeloma suggests that these malignancies may also represent promising targets for inhibitors of Ras signaling. While it is impossible to predict the outcome of the clinical trials, the biologic properties of these inhibitors are potentially informative because transformation-specific mechanisms are targeted.

Acknowledgment

We thank Dr Kristine A. Henningfeld for help with the figures and for critical reading of the manuscript.

References

- Spring SR. G protein mechanisms: insights from structural analysis. *Annu Rev Biochem*. 1997;66:639-678.
- Bos JL. Ali in the family? New insights and questions regarding interconnectivity of Ras, Rap1 and Ral. *EMBO J*. 1998;17:6776-6782.
- Robello A, Martínez CA. Ras proteins: recent advances and new functions. *Blood*. 1999;94:2971-2980.
- Pellis S, Divjak M, Romanowski P, et al. Developmentally-regulated expression of murine K-ras isoforms. *Oncogene*. 1997;15:1781-1788.
- Willinghofer A. Signal transduction via Ras. *Biol Chem*. 1998;379:933-937.
- Van Aelst L, White M, Wigler MH. Ras partners. *Cold Spring Harbor Symp Quant Biol*. 1994;59:181-186.
- Marshall CJ. Ras effectors. *Curr Opin Cell Biol*. 1996;8:197-204.
- Katz ME, McCormick F. Signal transduction from multiple Ras effectors. *Curr Opin Genet Dev*. 1997;7:75-79.
- Glassel JA, Farnsworth CC. Role of protein modification reactions in programming interactions between Ras-related GTPases and cell membranes. *Annu Rev Cell Biol*. 1994;10:181-205.
- Zhang FL, Casey PJ. Protein prenylation: molecular mechanisms and functional consequences. *Annu Rev Biochem*. 1996;65:241-269.
- Golt M. Protein prenylation, et cetera: signal transduction in two dimensions. *Science*. 1997;275:1750-1751.
- Mumby SM. Reversible palmitoylation of signaling proteins. *Curr Opin Cell Biol*. 1997;9:148-154.
- Casey PJ, Seabra MC. Protein prenyltransferases. *J Biol Chem*. 1996;271:5289-5292.
- Reiss Y, Goldstein JL, Seabra MC, Casey PJ, Brown MS. Inhibition of purified p21^{ras} farnesyl protein transferase by cys-AAAX tetrapeptides. *Cell*. 1990;62:81-88.
- Reiss Y, Stradley SJ, Gierach LM, Brown MS, Goldstein JL. Sequence requirement for peptide recognition by rat brain p21^{ras} protein farnesyltransferase. *Proc Natl Acad Sci U S A*. 1991;88:732-736.
- Yokoyama K, Goodwin GW, Ghomashchi F, Glassel JA, Gelb MH. A protein geranylgeranyltransferase from bovine brain: implications for protein prenylation specificity. *Proc Natl Acad Sci U S A*. 1991;88:5302-5306.
- Moore SL, Schaber MD, Moesser SD, et al. Sequence dependence of protein isoprenylation. *J Biol Chem*. 1991;266:14603-14610.
- Trueblood CE, Ohya Y, Rine J. Genetic evidence for in vivo cross-specificity of the CAAX-box protein prenyltransferases farnesyltransferase and geranylgeranyltransferase I in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 1993;13:4280-4275.
- Pollicena P, Scholten JD, Zimmerman K, Creswell M, Huang CC, Miller WT. Involvement of the alpha subunit of farnesyl-protein transferase in substrate recognition. *Biochemistry*. 1996;35:13494-13500.
- Trueblood CE, Boyartchuk VL, Rine J. Substrate specificity determinants in the farnesyltransferase beta-subunit. *Proc Natl Acad Sci U S A*. 1997;94:10774-10779.
- Park H-W, Bodulur SR, Moormaw JF, Casey PJ, Boese LS. Crystal structure of protein farnesyltransferase at 2.25 Angstrom resolution. *Science*. 1997;275:1800-1804.
- Akopyants TN, Couedel Y, Orlowski M, Fournie-Zalauki MC, Roques BP. Proteolytic processing of farnesylated peptides: assay and partial purification from pig brain membranes of an endopeptidase which has the characteristics of E.C.
- 3.4.24.15. *Biochem Biophys Res Commun*. 1994;198:787-794.
- Boyartchuk VL, Ashby MN, Rine J. Modulation of Ras and a-factor function by carboxy-terminal proteolysis. *Science*. 1997;275:1796-1800.
- Hancock J, Magee A, Childs J, Marshall C. All ras proteins are polyisoprenylated but only some are palmitoylated. *Cell*. 1989;57:1167-1177.
- Milligan G, Parenti M, Magee AI. The dynamic role of palmitoylation in signal transduction. *Trends Biochem Sci*. 1995;20:181-187.
- Ross EM. Palmitoylation in G-protein signaling pathways. *Curr Biol*. 1996;5:107-109.
- Dudler T, Gelb MH. Palmitoylation of Ha-Ras facilitates membrane binding, activation of downstream effectors and mitotic maturation in *Xenopus oocytes*. *J Biol Chem*. 1996;271:11541-11547.
- Liu L, Dudler T, Gelb MH. Purification of a protein palmitoyltransferase that acts on H-Ras protein and on a C-terminal N-Ras peptide. *J Biol Chem*. 1996;271:23269-23276.
- Camp LA, Verkruyse LA, Afendis SJ, Slaughter CA, Hofmann SL. Molecular cloning and expression of palmitoyl-protein thioesterase. *J Biol Chem*. 1994;269:23212-23219.
- Treisman R. Regulation of transcription by MAP kinase cascades. *Curr Opin Cell Biol*. 1996;8:205-215.
- Fanger GR, Gerwins P, Widmann C, Jarpe MB, Johnson GL. MEKs, GCKs, MLKs, PAKs, TAKs, and Tpl2: upstream regulators of the c-Jun amino-terminal kinases? *Curr Opin Genet Dev*. 1997;7:67-74.
- Robinson MJ, Cobb MH. Mitogen-activated protein kinase pathways. *Curr Opin Cell Biol*. 1997;9:180-188.
- Garrington TP, Johnson GL. Organization and regulation of mitogen-activated protein kinase signaling pathways. *Curr Opin Cell Biol*. 1999;11:211-218.
- Schaeffer HJ, Weber MJ. Mitogen-activated protein kinases: specific messages from ubiquitous messengers. *Mol Cell Biol*. 1999;19:2435-2444.
- Elion EA. Routing MAP kinase cascades. *Science*. 1998;281:1625-1628.
- Schlessinger J. How receptor tyrosine kinases activate Ras. *Trends Biol Sci*. 1993;18:273-275.
- Marshall CJ. Specificity of receptor tyrosine kinase signaling: transient versus sustained extracellular signal-regulated kinase activation. *Cell*. 1995;80:179-185.
- Marshall CJ. Raf gets it together. *Nature*. 1996;383:127-128.
- Porter AC, Vaillancourt RR. Tyrosine kinase receptor-activated signal transduction pathways which lead to oncogenesis. *Oncogene*. 1998;17:13434-13452.
- Pawson T, Saxton TM. Signaling networks—do all roads lead to the same genes? *Cell*. 1999;97:875-878.
- Adachi T, Alam R. The mechanism of IL-5 signal transduction. *Am J Physiol*. 1998;275:C623-C633.
- Guthridge MA, Stomski FC, Thomas D, et al. Mechanism of activation of the GM-CSF, IL-3, and IL-5 family of receptors. *Stem Cells*. 1998;16:301-313.
- Daum G, Eisenmann-Tappe I, Fries HW, Troppmair J, Rapp U. The ins and outs of Raf kinases. *Trends Biol Sci*. 1994;19:474-480.
- Callling AD, Schaeffer H-J, Reuter CWM, Reddy GR, Weber MJ. A proline-rich sequence unique to MEK1 and MEK2 is required for Raf binding and regulates MEK function. *Mol Cell Biol*. 1995;15:5214-5225.
- Reuter CWM, Callling AD, Jelinek T, Weber MJ. Biochemical analysis of MEK activation in NIH3T3 fibroblasts. *J Biol Chem*. 1995;270:7644-7655.
- Patriotic C, Makris A, Chernoff J, Tschlis PN. Tpl-2 acts in concert with Ras and Raf-1 to activate mitogen-activated protein kinase. *Proc Natl Acad Sci U S A*. 1994;91:9755-9759.
- Samaron A, Ahmad TB, Carlile GW, Pappin D, Narsimhan RP, Ley SC. Activation of MEK-1 and SEK-2 by Tpl-2 proto-oncoprotein, a novel MAP kinase kinase. *EMBO J*. 1996;15:817-826.
- Posada J, Yaw N, Ahn NG, Vande-Woude GF, Cooper JA. Mos stimulates MAP kinase in *Xenopus oocytes* and activates MAP kinase kinase in vitro. *Mol Cell Biol*. 1993;13:2546-2553.
- Bardwell L, Thorner J. A conserved motif at the amino termini of MEKs might mediate high affinity interaction with the cognate MAPKs. *Trends Biol Sci*. 1996;21:373-374.
- Crews CM, Alessandrini A, Erikson RL. The primary structure of MEK, a protein that phosphorylates the ERK gene product. *Science*. 1992;258:478-480.
- Wu J, Harrison JK, Dont P, Lynch KR, Weber MJ, Sturgill TW. Identification and characterization of a new mammalian mitogen-activated protein kinase kinase, MKK2. *Mol Cell Biol*. 1993;13:4539-4548.
- Zheng C-F, Guan K-L. Cloning and characterization of two distinct human extracellular signal-regulated kinase activator kinases, MEK1 and MEK2. *J Biol Chem*. 1993;268:11435-11439.
- Xing J, Ginty DD, Greenberg ME. Coupling of the Ras-MAPK pathway to gene activation by RSK2, a growth factor-regulated CREB kinase. *Science*. 1996;273:959-963.
- Jaaro H, Rubinfeld H, Hancock T, Seger R. Nuclear translocation of mitogen-activated protein kinase kinase (MEK1) in response to mitogenic stimulation. *Proc Natl Acad Sci U S A*. 1997;94:3742-3747.
- Felg LA, Urano T, Cantor S. Evidence for a Ras/Raf signaling cascade. *Trends Biochem Sci*. 1996;21:438-441.
- Carpenter CL, Cantley LC. Phosphoinositide kinases. *Curr Opin Cell Biol*. 1996;8:153-158.
- Graham SM, Cox AD, Drivas G, Rush MG, D'Eustachio P, Der CJ. Aberrant function of the Ras-related protein TC21/R-Ras2 triggers malignant transformation. *Mol Cell Biol*. 1994;14:4108-4115.
- Cox AD, Briva TR, Lowe DG, Der CJ. R-Ras induces malignant, but not morphologic, transformation of NIH3T3 cells. *Oncogene*. 1994;9:3281-3288.
- Quiliani LA, Castro AF, Rogers-Graham KS, Martin CB, Der CJ, Bl C. M-Ras/R-Ras3, a transforming ras protein regulated by SOS1, GRF1, and p120 Ras GTPase-activating protein, interacts with the putative Ras effector AF6. *J Biol Chem*. 1999;274:23850-23857.
- Bos JL. RAS oncogenes in human cancer: a review. *Cancer Res*. 1989;49:4682-4689.
- Clark GJ, Der CJ. Ras proto-oncogene activation in human malignancy. In: Garrett CT, Sul S, eds. *Cellular Cancer Markers*. Totowa, NJ: Humana Press; 1995:17-52.
- Janssen J, Steenvoorden A, Lyons J, et al. Ras gene mutations in acute and chronic myelocytic leukemias, chronic myeloproliferative disorders, and myelodysplastic syndromes. *Proc Natl Acad Sci U S A*. 1987;84:9228-9232.
- Bos JL, Verlaan-de-Vries M, van der Eb AJ, et al. Mutations in N-Ras predominate in acute myeloid leukemia. *Blood*. 1987;69:1237-1241.
- Farr CJ, Saiki RK, Erlich HA, McCormick F, Marshall CJ. Analysis of Ras gene mutations in acute myeloid leukemia by polymerase chain reaction

- and oligonucleotide probes. *Proc Natl Acad Sci U S A*. 1988;85:1629-1633.
65. Padua RA, Carter G, Hughes D, et al. Ras mutations in myelodysplasia detected by amplification, oligonucleotide hybridization and transformation. *Leukemia*. 1988;2:503-510.
 66. Senn HP, Tran-Thang C, Wodnar-Filipowicz A, et al. Mutational analysis of the N-RAS proto-oncogene in active and remission phase acute leukemias. *Int J Cancer*. 1988;41:59-64.
 67. Takkenz D, Farr CJ, Marshall CJ. Ras genes and acute myeloid leukemia. *Br J Haematol*. 1988;71:1-0.
 68. Browett PJ, Vaxley JC, Norton JD. Activation of Harvey ras oncogene by mutation at codon 12 is very rare in hematopoietic malignancies. *Leukemia*. 1989;3:86-88.
 69. Browett PJ, Norton JD. Analysis of RAS gene mutations and methylation state in human leukemias. *Oncogene*. 1989;4:1029-1036.
 70. Parker J, Mulli GJ. Ras and myelodysplasia: lessons from the last decade. *Semin Hematol*. 1996;33:206-224.
 71. Byrne JL, Marshall CJ. The molecular pathophysiology of myeloid leukemias: Ras revisited. *Br J Haematol*. 1998;100:256-264.
 72. Vogelstein B, Clinch CI, Preisler AC, et al. Ras gene mutations in childhood acute myeloid leukemia: a Pediatric Oncology Group study. *Genes Chromosomes Cancer*. 1990;2:159-162.
 73. Hirsch-Ginsberg C, LeMaistre AC, Kantarjian H, et al. Ras mutations are rare events in Philadelphia chromosome-negative/bcr gene rearrangement-negative chronic myelogenous leukemia, but are prevalent in chronic myelomonocytic leukemia. *Blood*. 1990;76:1214-1219.
 74. Hnilok M, Laif Bergsagel P, Anderson KC. Multiple myeloma: Increasing evidence for a multi-step transformation process. *Blood*. 1998;91:3-21.
 75. Nerl A, Knowles DM, Greco A, McCormick F, Datta-Favara R. Analysis of Ras oncogene mutations in human lymphoid malignancies. *Proc Natl Acad Sci U S A*. 1988;85:9260-9272.
 76. Nerl A, Murphy JP, Cro L, et al. Ras oncogene mutation in multiple myeloma. *J Exp Med*. 1989;170:1716-1725.
 77. Tanaka K, Takeuchi M, Asakura H, Dohy H, Kamada N. A high frequency of N-Ras oncogene mutations in multiple myeloma. *Int J Hematol*. 1992;56:119-127.
 78. Corralini P, Ladetto M, Voena C, et al. Mutational activation of N- and K-RAS oncogenes in plasma cell dyscrasias. *Blood*. 1993;81:2708-2713.
 79. Hunter T. Oncoprotein networks. *Cell*. 1997;88:333-348.
 80. Snwyers CL, Denny CT. Chronic myelomonocytic leukemia: Tel-3-kinase what Ets all about. *Cell*. 1994;77:171-173.
 81. Tobai K, Pagliuca A, Bhatt B, Bailey N, Layton DM, Mulli GJ. Mutation of the human FMS gene (M-CSF receptor) in myelodysplastic syndromes and acute myeloid leukemia. *Leukemia*. 1990;4:486-489.
 82. Portua RA, Guinn BA, Al-Sabah AI, et al. Ras, FMS and p53 mutations and poor clinical outcome in myelodysplasias: a 10-year follow-up. *Leukemia*. 1998;12:887-892.
 83. Nakata Y, Kimura A, Katoh O, et al. c-kit point mutation of extracellular domain in patients with myeloproliferative disorders. *Br J Haematol*. 1995;91:661-663.
 84. Bulthor C, Honz BM, Walker P, Sepp NT, Grabbe J. Identification of activating c-kit mutations in adult, but not in childhood-onset indolent mastocytosis: a possible explanation for divergent clinical behavior. *J Invest Dermatol*. 1998;111:1227-1231.
 85. Nagata H, Worobec AS, Oh CK, et al. Identification of a point mutation in the catalytic domain of the protooncogene c-kit in peripheral blood mononuclear cells of patients who have mastocytosis with an associated hematologic disorder. *Proc Natl Acad Sci U S A*. 1995;92:10560-10564.
 86. Kiyoi H, Naoe T, Nakano Y, et al. Prognostic implication of FLT3 and N-Ras gene mutations in acute myeloid leukemia. *Blood*. 1999;93:3074-3080.
 87. Dosi M, Wang S, Lemischka IR. Mitogenic signaling and substrate specificity of the FIK2/Flt3 receptor tyrosine kinase in fibroblasts and interleukin 3-dependent hematopoietic cells. *Mol Cell Biol*. 1993;13:6572-6585.
 88. Rohrschneider LR, Bourette RP, Lioubin MN, Aigle PA, Myles GM, Carlberg K. Growth and differentiation signals regulated by the M-CSF receptor. *Mol Reprod Dev*. 1997;46:96-103.
 89. Elmberger PG, Lozano MD, Weisenburger DD, Sanger W, Chan WC. Transcripts of the *nm-alk* fusion gene in anaplastic large cell lymphoma, Hodgkin's disease, and reactive lymphoid lesions. *Blood*. 1995;86:3517-3521.
 90. Waggott W, Lo YM, Bastard C, et al. Detection of NPM-ALK DNA rearrangement in CD30 positive anaplastic large cell lymphoma. *Br J Haematol*. 1995;89:905-907.
 91. Golub T, Barker G, Lovett M, Gilliland D. Fusion of PDGF receptor β to a novel ets-like gene, *tel*, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell*. 1994;77:307-316.
 92. Jousset C, Carron C, Bouraux A, et al. A domain of TEL conserved in a subset of ETS proteins defines a specific oligomerization interface essential to the mitogenic properties of the TEL-PDGFR β oncoprotein. *EMBO J*. 1997;16:69-82.
 93. Papadopoulos P, Ridge SA, Boucher CA, Stocking C, Wiedemann LM. The novel activation of *abl* by fusion to an ets-related gene, *tel*. *Cancer Res*. 1995;55:34-38.
 94. Golub TR, Goga A, Barker GF, et al. Oligomerization of the ABL tyrosine kinase by the Ets protein TEL in human leukemia. *Mol Cell Biol*. 1996;16:4107-4116.
 95. Kurzrock R, Guterman J, Talpaz M. The molecular genetics of Philadelphia chromosome-positive leukemias. *N Engl J Med*. 1988;319:990-998.
 96. Faderl S, Talpaz M, Estrow Z, O'Brien S, Kurzrock R, Kantarjian HM. The biology of chronic myeloid leukemia. *N Engl J Med*. 1999;341:164-172.
 97. Zou X, Calame K. Signaling pathways activated by oncogenic forms of Abl tyrosine kinase. *J Biol Chem*. 1999;274:18141-18144.
 98. Xu G, O'Connell P, Viskochil D, et al. The neurofibromatosis type 1 gene encodes a protein related to GAP. *Cell*. 1990;62:599-608.
 99. Niemeyer CM, Arico M, Basso G, et al. Chronic myelomonocytic leukemia in childhood: a retrospective analysis of 110 cases. *Blood*. 1997;89:3534-3543.
 100. Shannon KM, O'Connell P, Martin GA, et al. Loss of the normal NF-1 allele from the bone marrow of children with type 1 neurofibromatosis and malignant myeloid disorders. *N Engl J Med*. 1994;330:597-601.
 101. Stiller CA, Chessells JM, Fitchett M. Neurofibromatosis and childhood leukemia/lymphoma: a population based UKCCSG study. *Br J Cancer*. 1994;70:969-972.
 102. Side L, Taylor B, Cayouette M, et al. Homozygous inactivation of the NF-1 gene in bone marrow cells from children with neurofibromatosis type 1 and malignant myeloid disorders. *N Engl J Med*. 1997;336:1713-1720.
 103. DeClue JE, Papageorge AG, Fletcher JA, et al. Abnormal regulation of mammalian p21ras contributes to malignant tumor growth in von Recklinghausen (type 1) neurofibromatosis. *Cell*. 1992;69:265-273.
 104. Kolra R, Paderanga DC, Olson K, Shannon KM. Genetic analysis is consistent with the hypothesis that NF-1 limits myeloid cell growth through p21ras. *Blood*. 1994;84:3435-3439.
 105. Bollag G, Clapp DW, Shih S, et al. Loss of NF-1 results in activation of the Ras signaling pathway and leads to aberrant growth in hematopoietic cells. *Nat Genet*. 1996;12:144-148.
 106. Largaespada DA, Brannan CI, Jenkins NA, Copeland NG. NF-1 deficiency causes Ras-mediated granulocyte/macrophage colony stimulating factor hypersensitivity and chronic myeloid leukemia. *Nat Genet*. 1996;12:137-143.
 107. Miyachi J, Asada M, Sasaki M, Tsunenitsu Y, Kojima S, Mizutani S. Mutations of N-ras gene in juvenile chronic myelogenous leukemia. *Blood*. 1994;83:2248-2254.
 108. Birnbaum RA, O'Marcalgh A, Wardak Z. NF1 and GM-CSF interact in myeloid leukemogenesis. *Mol Cell*. 2000;5:189-195.
 109. MacKenzie KL, Dolnikov A, Millington M, Shouan Y, Symonds G. Mutant N-ras induces myeloproliferative disorders and apoptosis in bone marrow repopulated mice. *Blood*. 1999;93:2043-2050.
 110. Saison-Behomcaras T, Torque B, Roy J, Chassignol M, Thuong NT, Helene C. Short modified antisense oligonucleotides directed against Ha-RAS point mutation induce selective cleavage of the mRNA and inhibit T24 cells proliferation. *EMBO J*. 1991;10:1111-1118.
 111. Mukhopadhyay T, Tolinsky M, Cavender AC, Ruth JA. Specific inhibition of K-ras expression and tumorigenicity of lung cancer cells by antisense RNA. *Cancer Res*. 1991;51:1744-1748.
 112. Shirogawa S, Furuse M, Yokoyama N, Sasazuki T. Altered growth of human colon cancer cell lines disrupted at activated Ki-Ras. *Science*. 1993;260:85-88.
 113. Kashani-Sabet M, Funato T, Florones VA, Fodstad O, Scanlon KJ. Suppression of the neoplastic phenotype in vivo by an anti-ras ribozyme. *Cancer Res*. 1994;54:900-902.
 114. Gibbs JB. Ras C-terminal processing enzymes: new drug targets? *Cell*. 1991;65:1-4.
 115. Tamarai F. Inhibitors of Ras farnesyltransferases. *Trends Biochem Sci*. 1993;18:349-353.
 116. Gibbs JB, Oliff A. Pharmaceutical research in molecular oncology. *Cell*. 1994;79:193-198.
 117. Gibbs JB, Oliff A, Kohl NE. Farnesyltransferase inhibitors: Ras research yields a potential cancer therapeutic. *Cell*. 1994;77:175-178.
 118. Lowy DR, Witkensen BM. Rational cancer therapy. *Nat Med*. 1995;1:747-748.
 119. Gibbs JB, Oliff A. The potential of farnesyltransferase inhibitors as cancer chemotherapeutics. *Annu Rev Pharmacol Toxicol*. 1997;37:143-166.
 120. Omer CA, Kohl NE. CA1A2X-competitive inhibitors of farnesyltransferase as anti-cancer agents. *Trends Pharmacol Sci*. 1997;18:437-444.
 121. Heinbrook DC, Oliff A. Therapeutic intervention and signaling. *Curr Biol*. 1998;10:284-288.
 122. Kato K, Cox AD, Hisaka MM, Graham SM, Buss JE, Der CJ. Isoprenoid addition to Ras protein is the critical modification for its membrane association and transforming activity. *Proc Natl Acad Sci U S A*. 1992;89:6403-6407.
 123. Kang MS, Stomerick DM, Zwolsen JH, Harry BS, Sunkara PS, Harrison BL. Farnesyl-derived inhibitors of Ras farnesyl transferase. *Biochem Biophys Res Commun*. 1995;217:245-249.
 124. Yonemoto M, Satoh T, Arakawa H, et al. J-104,871, a novel farnesyltransferase inhibitor, blocks Ras farnesylation in vivo in a farnesyl pyrophosphate-competitive manner. *Mol Pharmacol*. 1998;54:1-7.
 125. James GL, Goldstein JL, Brown MS, et al. Benzodiazepine peptidomimetics: potent inhibitors of Ras farnesylation in animal cells. *Science*. 1993;260:1937-1942.

1668 REUTER et al

BLOOD, 1 SEPTEMBER 2000 • VOLUME 96, NUMBER 5

126. Jaino GL, Goldstein JL, Brown MS. Polytyrosine and CVIM sequences of K-RasB dictate specificity of prenylation and confer resistance to benzodiazepine peptidomimetic in vitro. *J Biol Chem*. 1995;270:6221-6226.
127. Dalton MD, Fante KS, Bechtold HA, et al. The farnesyl protein transferase inhibitor BZA-5B blocks farnesylation of nuclear lamins and p21ras but does not affect their function or localization. *Cancer Res*. 1995;55:3295-3304.
128. Kohl NE, Mosser SD, DeSolms SJ, et al. Selective inhibition of ras-dependent transformation by a farnesyltransferase inhibitor. *Science*. 1993; 260:1934-1937.
129. Kohl NE, Wilson FR, Mosser SD, et al. Protein farnesyltransferase inhibitors block the growth of ras-transformed tumors in nude mice. *Proc Natl Acad Sci U S A*. 1994;91:9141-9145.
130. Prendergast GC, Davide JP, DeSolms SJ, et al. Farnesyltransferase inhibition causes morphological reversion of ras-transformed cells by a complex mechanism that involves regulation of the actin cytoskeleton. *Mol Cell Biol*. 1994;14: 4193-4202.
131. Lebowitz PF, Sakamuro D, Prendergast GC. Farnesyl transferase inhibitors induce apoptosis of Ras-transformed cells denied subcellular attachment. *Cancer Res*. 1997;57:708-713.
132. Emanuel PD, Snyder RC, Willey T, Gopuram B, Castiberry RP. Inhibition of juvenile myelomonocytic leukemia cell growth in vitro by farnesyltransferase inhibitors. *Blood*. 2000;95:839-845.
133. Koltan KS, Culbertson JC, DeSolms SJ, et al. NMR studies of novel inhibitors bound to farnesyl-protein transferase. *Protein Sci*. 1995;4:681-688.
134. Barrington RE, Subler MA, Rands E, et al. A farnesyltransferase inhibitor induces tumor regression in transgenic mice harboring multiple oncogenic mutations by mediating alterations in both cell cycle control and apoptosis. *Mol Cell Biol*. 1998;18:85-92.
135. Kohl NE, Omer CA, Conner MW, et al. Inhibition of farnesyltransferase induces regression of mammary and salivary carcinomas in ras transgenic mice. *Nat Med*. 1995;1:782-787.
136. Sepp-Loranzino L, Ma Z, Rands E, et al. A peptidomimetic inhibitor of farnesyl protein transferase blocks the anchorage-dependent and -independent growth of human tumor cell lines. *Cancer Res*. 1995;55:5302-5309.
137. Mangues R, Corral T, Kohl NE, et al. Antitumor effect of a farnesyl protein transferase inhibitor in mammary and lymphoid tumors overexpressing N-RAS in transgenic mice. *Cancer Res*. 1998;58: 1253-1259.
138. Cox AD, Garcia AM, Westwick JK, et al. The CAAX peptidomimetic compound B581 specifically blocks farnesylation, but not geranylgeranylation or myristylation, oncogenic Ras signaling and transformation. *J Biol Chem*. 1994;269:19203-19206.
139. Qian Y, Blaskovich MA, Saleem M, et al. Design and structural requirements of potent peptidomimetic inhibitors of p21ras farnesyltransferase. *J Biol Chem*. 1994;269:12410-12413.
140. Sun J, Qian Y, Hamilton AD, Sebti SM. Ras CAAX peptidomimetic FTI-276 selectively blocks tumor growth in nude mice of a human lung carcinoma with K-Ras mutation and p53 deletion. *Cancer Res*. 1995;55:4243-4247.
141. Lerner EC, Qian Y, Blaskovich MA, et al. Ras CAAX peptidomimetic FTI-277 selectively blocks oncogenic Ras signaling by inducing cytoplasmic accumulation of inactive Ras-Raf complexes. *J Biol Chem*. 1995;270:26802-26806.
142. Bradel M, Pullack IF, Freund JM, Hamilton AD, Sebti SM. Inhibition of Ras and related G-proteins as a therapeutic strategy for blocking malignant glioma growth. *Neurosurgery*. 1998;43:124-131.
143. Bornhard EJ, McKenna WG, Hamilton AD, et al. Inhibiting Ras prenylation increases the radiosensitivity of human tumor cell lines with activating mutations of ras oncogenes. *Cancer Res*. 1998; 58:1754-1761.
144. Nagasu T, Yoshimatsu K, Rowell C, Lewis MD, Garcia AM. Inhibition of human tumor xenograft growth by treatment with the farnesyl transferase inhibitor B958. *Cancer Res*. 1995;55:5310-5314.
145. Bishop WR, Bond R, Petrin J, et al. Novel tricyclic inhibitors of farnesyl protein transferase. *J Biol Chem*. 1995;270:30611-30618.
146. Njoroge FG, Vibulbhan B, Pinto P, et al. Potent, selective, and orally bioavailable tricyclic pyridyl acetamide N-oxide inhibitors of farnesyl protein transferase with enhanced in vivo antitumor activity. *J Med Chem*. 1998;41:1561-1567.
147. Njoroge FG, Taveras AG, Kelly J, et al. (1-4-[2-[4-(8-chloro-3,10-dibromo-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2b]pyridin-11(R)-yl]-1-piperidinyl)-2-oxo-ethyl]-1-piperidinecarboxamide (SCH66336): a very potent farnesyl protein transferase inhibitor as a novel antitumor agent. *J Med Chem*. 1998;41:4890-4902.
148. Mallams AK, Rossman RR, Doll RJ, et al. Inhibitors of farnesyl protein transferase. 4-Amido, 4-carbamoyl, and 4-carboxamido derivatives of 1-(8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2b]pyridin-11-yl)piperazine and 1-(3-bromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2b]pyridin-11-yl)piperazine. *J Med Chem*. 1998;41:877-893.
149. Liu M, Bryant MS, Chen J, et al. Effects of SCH59228, an orally bioavailable farnesyl protein transferase inhibitor, on the growth of oncogene-transformed fibroblasts and a human colon carcinoma xenograft in nude mice. *Cancer Chemother Pharmacol*. 1999;43:50-58.
150. Patel DV, Gordon EM, Schmidt RJ, et al. Phosphoryl acid-based bisubstrate analog inhibitors of Ras farnesyl protein transferase. *J Med Chem*. 1995;38:435-442.
151. Manne V, Yan N, Carboni JM, et al. Bisubstrate inhibitors of farnesyltransferase: a novel class of specific inhibitors of ras-transformed cells. *Oncogene*. 1995;10:1763-1779.
152. Patel DV, Young MG, Robinson SP, Hunihan L, Dean BJ, Gordon EM. Hydroxamic acid-based bisubstrate analog inhibitors of Ras farnesyl protein transferase. *J Med Chem*. 1996;39:4197-4210.
153. Geib MH, Tamanol F, Yokoyama K, Ghomashchi F, Esson K, Gould MN. The inhibition of protein prenyltransferases by oxygenated metabolites of limonene and perillyl alcohol. *Cancer Lett*. 1995; 91:169-175.
154. Hara M, Akasaka K, Akinaga S, et al. Identification of Ras farnesyltransferase inhibitors by microbial screening. *Proc Natl Acad Sci U S A*. 1993;90:2281-2285.
155. Nagase T, Kawata S, Tamura S, et al. Manumycin and gilotoxin derivative KT7595 block Ras farnesylation and cell growth but do not disturb lamin farnesylation and localization in human tumor cells. *Br J Cancer*. 1997;76:1001-1010.
156. Kainuma O, Asano T, Hasegawa M, et al. Inhibition of growth and invasive activity of human pancreatic cancer cells by a farnesyltransferase inhibitor, manumycin, Pancreas. 1997;15:379-383.
157. Jayasuriya H, Ball RG, Zink DL, et al. Barcelona acid A, a new farnesyl-protein transferase inhibitor from *Phoma* species. *J Nat Prod*. 1995;58: 986-991.
158. Van der Pyl D, Cans P, Debernard JJ, et al. RPR113228, a novel farnesyl-protein transferase inhibitor produced by *Chrysosporium lobatum*. *J Antibiot (Tokyo)*. 1995;48:736-737.
159. Silverman KC, Cascales C, Genilloud O, et al. Actinoplastic acids A and B as novel inhibitors of farnesyl-protein transferase. *Appl Microbiol Biotechnol*. 1995;43:810-816.
160. Silverman KC, Jayasuriya H, Cascales C, et al. Organic acid, a potent inhibitor of Ras farnesyl-protein transferase. *Biochem Biophys Res Commun*. 1997;232:478-481.
161. Sturm S, Gil RR, Chai HB, et al. Lupane derivatives from *Lophopetalum wallichii* with farnesyl protein transferase inhibitory activity. *J Nat Prod*. 1990;59:658-663.
162. Sekizawa R, Iinuma H, Naganawa H, et al. Isolation of novel saquayamycins as inhibitors of farnesyl-protein transferase. *J Antibiot (Tokyo)*. 1996;49:487-490.
163. Tsuda M, Muraoka Y, Takeuchi T, Sekizawa R, Umezawa K. Stereospecific synthesis of a novel farnesyl protein transferase inhibitor, valinocin and its analogues. *J Antibiot (Tokyo)*. 1996;49: 1031-1035.
164. Lee S, Park S, Oh JW, Yang C. Natural inhibitors for protein prenyltransferase. *Planta Med*. 1998; 64:303-308.
165. Gibbs JB, Porriano DL, Mosser SD, et al. Selective inhibition of farnesyl-protein transferase blocks Ras processing in vivo. *J Biol Chem*. 1993; 268:7617-7620.
166. James GL, Brown MS, Cobb MH, Goldstein JL. Benzodiazepine peptidomimetic BZA-5B interrupts the MAP kinase activation pathway in H-Ras-transformed Rat-1 cells, but not in untransformed cells. *J Biol Chem*. 1994;269:27705-27714.
167. James G, Goldstein JL, Brown MS. Resistance of K-Ras^{B12} proteins to farnesyltransferase inhibitors in Rat1 cells. *Proc Natl Acad Sci U S A*. 1990; 87:4454-4458.
168. Sun J, Qian Y, Hamilton AD, Sebti SM. Both farnesyltransferase and geranylgeranyltransferase I inhibitors are required for inhibition of oncogenic K-Ras prenylation but each alone is sufficient to suppress human tumor growth in nude mouse xenografts. *Oncogene*. 1998;16:1467-1473.
169. Whyte DB, Kirschmeier P, Hockenberry TN, et al. K-Ras and N-Ras are geranylgeranylated in cells treated with farnesyl protein transferase inhibitors. *J Biol Chem*. 1997;272:14459-14464.
170. Zhang FL, Kirschmeier P, Carr D, et al. Characterization of Ha-Ras, N-Ras, Ki-Ras4A, and Ki-Ras4B as in vivo substrates for farnesyl protein transferase and geranylgeranyl protein transferase type I. *J Biol Chem*. 1997;272:10232-10239.
171. Rowell CA, Kowalczyk JJ, Lewis MD, Garcia AM. Direct demonstration of geranylgeranylation and farnesylation of Ki-Ras in vivo. *J Biol Chem*. 1997;272:14093-14097.
172. Lerner EC, Zhang TT, Knowles DB, Qian Y, Hamilton AD, Sebti SM. Inhibition of the prenylation of K-RAS, but not H- or N-RAS is highly resistant to CAAX peptidomimetics and requires both a farnesyltransferase and a geranylgeranyltransferase I inhibitor in human tumor cell lines. *Oncogene*. 1997;15:1283-1288.
173. Lerner EC, Qian Y, Hamilton AD, Sebti SM. Disruption of oncogenic K-Ras4B processing and signaling by a potent geranylgeranyltransferase I inhibitor. *J Biol Chem*. 1995;270:26770-26773.
174. Lebowitz PF, Davide JP, Prendergast GC. Evidence that farnesyltransferase inhibitors suppress Ras transformation by interfering with Rho activity. *Mol Cell Biol*. 1995;15:6613-6622.
175. Lebowitz PF, Prendergast GC. Non-Ras targets of farnesyltransferase inhibitors: focus on Rho. *Oncogene*. 1998;17:1439-1445.
176. Booden MA, Baker TL, Sokki PA, Dar CJ, Punko SG, Buss JE. A non-farnesylated Ha-Ras protein can be palmitoylated and trigger potent differentiation and transformation. *J Biol Chem*. 1999; 274:1423-1431.
177. Cox AD, Der CJ. Farnesyltransferase inhibitors and cancer treatment: targeting simply Ras? *Biochim Biophys Acta*. 1997;1333:F61-F71.
178. Du W, Lebowitz PF, Prendergast GC. Cell growth

- inhibition by farnesyltransferase inhibitors is mediated by gain of geranylgeranylated RhoB. *Mol Cell Biol*. 1999;19:1831-1840.
179. Adamson P, Marshall CJ, Hall A, Tilbrook PA. Post-translational modifications of p21/rho proteins. *J Biol Chem*. 1992;267:20033-20038.
 180. Sepp-Lorenzino L, Rosen N. A farnesyl-protein transferase inhibitor induces p21 expression and G1 block in p53 wild type tumor cells. *J Biol Chem*. 1998;273:20243-20251.
 181. Hung WC, Chuang LY. Involvement of caspase family proteases in FPI inhibitor II-induced apoptosis in human ovarian cancer cells. *Int J Cancer*. 1998;12:1339-1342.
 182. Hung WC, Chuang LY. The farnesyltransferase inhibitor, FPI inhibitor II upregulates Bax and Bcl-2 expression and induces apoptosis in human ovarian cancer cells. *Int J Oncol*. 1998;12:137-140.
 183. Norgaard P, Law B, Joseph H, et al. Treatment with farnesyl-protein transferase inhibitor induces regression of mammary tumors in transforming growth factor (TGF) alpha and TGF alpha/neu transgenic mice by inhibition of mitogenic activity and induction of apoptosis. *Clin Cancer Res*. 1999;5:35-42.
 184. Moasser MM, Sepp-Lorenzino L, Kohl NE, Oliff A, Balog A, Su DS. Farnesyl transferase inhibitors cause enhanced mitotic sensitivity to taxol and opothilons. *Proc Natl Acad Sci U S A*. 1998;95:1369-1374.
 185. DeSolms SJ, Giuliani EA, Graham St., et al. N-Arylalkyl pseudopeptide inhibitors of farnesyltransferase. *J Med Chem*. 1998;41:2651-2658.
 186. McNamara DJ, Dobrusin E, Leonard DM, et al. C-terminal modifications of histidyl-N-benzylglycyl derivatives to give improved inhibition of Ras farnesyltransferase, cellular activity, and anticancer activity in mice. *J Med Chem*. 1997;40:3319-3322.
 187. Leftleris K, Kline T, Vile GD, et al. Development of potent inhibitors of Ras farnesyltransferase possessing cellular and in vivo activity. *J Med Chem*. 1998;39:224-236.
 188. Jansen B, Schlagbauer-Wadl H, Kahr H, et al. Novel Ras antagonist blocks human melanoma growth. *Proc Natl Acad Sci U S A*. 1999;96:14010-14024.
 189. Brown MS, Goldstein JL, Paris KJ, Bunler JP, Marsters JC Jr. Tetrapeptide inhibitors of protein farnesyltransferase: amino-terminal substitution in phenylalanine-containing tetrapeptides restores farnesylation. *Proc Natl Acad Sci U S A*. 1992;89:8313-8316.
 190. Mahgoub N, Taylor BR, Gratton M, et al. In vitro and in vivo effects of a farnesyltransferase inhibitor on NF-1-deficient hematopoietic cells. *Blood*. 1999;94:2469-2476.
 191. Ito T, Kawata S, Tamura S, et al. Suppression of human pancreatic cancer growth in BALB/c nude mice by monomycin, a farnesyl-protein transferase inhibitor. *Jpn J Cancer Res*. 1996;87:113-116.
 192. Liu M, Bryant MS, Chen J, et al. Antitumor activity of SCH66336, an orally bioavailable tricyclic inhibitor of farnesyl protein transferase, in human tumor xenograft models and Wap-ras transgenic mice. *Cancer Res*. 1998;58:4947-4956.
 193. Prendergast GC, Davide JP, Lobowitz PF, Wachster-Roya R, Kohl NE. Resistance of a variant ras-transformed cell line to phenotypic reversion by farnesyl transferase inhibitors. *Cancer Res*. 1996;56:2626-2632.
 194. Del Villar K, Urano J, Guo L, Tamanoi F. A mutant form of human protein farnesyltransferase exhibits increased resistance to farnesyltransferase inhibitors. *J Biol Chem*. 1999;274:27010-27017.
 195. Qian Y, Vogt A, Vasudevan A, Sefti SM, Hamilton AD. Selective inhibition of type-I geranylgeranyltransferase in vitro and in whole cells by CAAL peptidomimetics. *Bioorg Med Chem*. 1998;6:293-299.
 196. Vogt A, Qian Y, McGuire TF, Hamilton AD, Sefti SM. Protein geranylgeranylation, not farnesylation, is required for the G1 to S phase transition in mouse fibroblasts. *Oncogene*. 1996;13:1991-1999.
 197. McGuire TF, Qian Y, Vogt A, Hamilton AD, Sefti SM. Platelet-derived growth factor receptor tyrosine phosphorylation requires protein geranylgeranylation but not farnesylation. *J Biol Chem*. 1996;271:27402-27407.
 198. Miquel K, Pradines A, Sun J, et al. GGT-298 induces G0-G1 block and apoptosis whereas FPI-277 causes G2-M enrichment in A549 cells. *Cancer Res*. 1997;57:1846-1850.
 199. Vogt A, Sun J, Qian Y, Hamilton AD, Sefti SM. The geranylgeranyltransferase-I inhibitor GGT-298 arrests human tumor cells in G0/G1 and induces p21(WAF1/CIP1/SDI1) in a p53-independent manner. *J Biol Chem*. 1997;272:27224-27229.
 200. Lantry LE, Zhang Z, Yao R, et al. Effect of farnesyltransferase inhibitor FPI-278 on established lung adenomas from A/J mice induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone. *Carcinogenesis*. 2000;21:113-116.
 201. Volker C, Miller RA, McCleary WR, et al. Effects of farnesylcysteine analogs on protein carboxyl methylation and signal transduction. *J Biol Chem*. 1991;266:21515-21522.
 202. Marciano D, Ben-Baruch G, Marom M, Egozi Y, Haklai R, Kloog Y. Farnesyl derivatives of rigid carboxylic acids—Inhibitors of Ras-dependent cell growth. *J Med Chem*. 1995;38:1267-1272.
 203. Marom M, Haklai R, Ben-Baruch G, Marciano D, Egozi Y, Kloog Y. Selective inhibition of Ras-dependent cell growth by farnesylthiosalicylic acid. *J Biol Chem*. 1995;270:22263-22270.
 204. Gana-Weisz M, Haklai R, Marciano D, Egozi Y, Ben-Baruch G, Kloog Y. The Ras antagonist S-farnesylthiosalicylic acid induces inhibition of MAPK activation. *Biochem Biophys Res Commun*. 1997;239:900-904.
 205. Haklai R, Weisz MG, Elad G, et al. Dislodgment and accelerated degradation of Ras. *Biochemistry*. 1998;37:1306-1314.
 206. Perez-Sala D, Gilbert BA, Rando RR, Canada FJ. Analogs of farnesylcysteine induce apoptosis in HL-60 cells. *FEBS Lett*. 1998;426:319-324.
 207. Aharonson Z, Gana-Weisz M, Varsano T, Haklai R, Mardano D, Kloog Y. Stringent structural requirements for anti-Ras activity of S-prenyl analogues. *Biochim Biophys Acta*. 1998;1406:40-50.
 208. Kadam S, McAlpine JB. Dorriginols: novel antifungal antibiotics that change the morphology of ras-transformed NIH3T3 cells to that of normal cells. III. Biological properties and mechanism of action. *J Antibiot (Tokyo)*. 1994;47:875-880.
 209. Chen Y. Selective inhibition of ras-transformed cell growth by a novel fatty acid-based chloromethyl ketone designed to target Ras endoprotease. *Ann N Y Acad Sci*. 1999;886:103-108.
 210. Alessi DR, Cuenda A, Cohen P, Dudley DT, Saltiel AD. PD 098059 is a specific inhibitor of the activation of mitogen-activated protein kinase kinase in vitro and in vivo. *J Biol Chem*. 1995;270:27489-27494.
 211. Dudley DT, Pang L, Decker SJ, Bridges AJ, Saltiel AR. A synthetic inhibitor of the mitogen-activated protein kinase cascade. *Proc Natl Acad Sci U S A*. 1995;92:7686-7689.
 212. DeSilva DR, Jones EA, Favata MF, et al. Inhibition of mitogen-activated protein kinase kinase blocks T cell proliferation but does not induce or prevent energy. *J Immunol*. 1998;160:4175-4181.
 213. Favata MF, Horiuchi KJ, Manos EJ, et al. Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J Biol Chem*. 1998;273:18623-18632.
 214. Williams DH, Wilkinson SE, Purton T, Lamont A, Flotow H, Murray EJ. Ro 09-2210 exhibits potent anti-proliferative effects on activated T-cells by selectively blocking MKK activity. *Biochemistry*. 1998;37:9579-9585.
 215. Cox LY, Molz J, Troll W, Garte SJ. Antipain-induced suppression of oncogene expression in H-ras-transformed NIH3T3 cells. *Cancer Res*. 1991;51:4810-4814.
 216. Itoh O, Kuroiwa S, Aisumi S, Umezawa K, Takeuchi T, Mar M. Induction by guanosine analogue oxanosine of reversion toward the normal phenotype of K-ras transformed rat kidney cells. *Cancer Res*. 1989;49:996-1000.
 217. Shindo-Okado N, Makabe O, Nagahara H, Nishimura S. Permanent conversion of mouse and human cells transformed by activated ras or raf genes to apparently normal cells by treatment with the antibiotic azalcyrosine. *Mol Carcinog*. 1989;2:159-167.
 218. Kumar CC, Prorock-Rogers C, Kelly J, et al. SCH51344 inhibits ras transformation by a novel mechanism. *Cancer Res*. 1995;55:5100-5117.
 219. Walsh AB, Dhanasekaran M, Bar-Sagi D, Kumar CC. SCH51344-induced reversal of Ras-transformation is accompanied by the specific inhibition of the Ras and Rac-dependent cell morphology pathway. *Oncogene*. 1997;15:2553-2560.
 220. Giardello FM, Hamilton SR, Krush AJ, et al. Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *N Engl J Med*. 1993;328:1313-1316.
 221. Verhuel HM, Panigrahy D, Yuan J, D'Amato RJ. Combination oral antiangiogenic therapy with thalidomide and sulindac inhibits tumour growth in rabbits. *Br J Cancer*. 1999;79:114-118.
 222. Herrmann C, Block C, Goison C, et al. Sulindac sulfide inhibits Ras signaling. *Oncogene*. 1998;17:1769-1778.
 223. Schulte TW, Blagosklonny MV, Romanova L, et al. Destabilization of Raf-1 by geldanamycin leads to disruption of the Raf-1-MEK-mitogen-activated protein kinase signaling pathway. *Mol Cell Biol*. 1996;16:5839-5845.
 224. Stebbins CE, Russo AA, Schelder C, Rosen N, Hart FU, Pavletich NP. Crystal structure of an HSP90-geldanamycin complex: targeting of a protein chaperone by an antitumor agent. *Cell*. 1997;89:239-250.
 225. Koera K, Nakamura K, Nakao K, et al. K-ras is essential for the development of the mouse embryo. *Oncogene*. 1997;15:1151-1159.
 226. Johnson L, Greenbaum D, Cichowski K, et al. K-ras is an essential gene in the mouse with partial functional overlap with N-ras. *Genes Dev*. 1997;11:2468-2481.
 227. Ulanoff H, Edelman W, Pellicer A, Kucherlapati R. The murine N-ras gene is not essential for growth and development. *Proc Natl Acad Sci U S A*. 1995;92:1709-1713.
 228. Casey S, Dautry F. Inactivation of the murine N-ras gene by gene targeting. *Oncogene*. 1992;12:2525-2528.
 229. Qiu RG, Chen J, McCormick F, Symons M. Role for Rho in Ras transformation. *Proc Natl Acad Sci U S A*. 1995;92:11781-11785.
 230. Khosravi-Far R, Solaki PA, Clark GJ, Kinch MS, Der CJ. Activation of Rac1, RhoA and mitogen-activated protein kinases is required for Ras transformation. *Mol Cell Biol*. 1995;15:6443-6453.
 231. Khosravi-Far R, White MA, Westwick JK, et al. Oncogenic Ras activation of Raf/mitogen-activated protein kinase-independent pathways is sufficient to cause tumorigenic transformation. *Mol Cell Biol*. 1996;16:3923-3933.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.